

What Is Adenomyosis? Unrecognized Condition Impacting Women's Lives. Affects Up to 1 in 5 Women

By [Dr. Joseph Mercola](#)

Theme: [Science and Medicine](#)

Global Research, June 26, 2024

[Mercola](#)

All Global Research articles can be read in 51 languages by activating the Translate Website button below the author's name (only available in desktop version).

To receive Global Research's Daily Newsletter (selected articles), [click here](#).

Click the share button above to email/forward this article to your friends and colleagues. Follow us on [Instagram](#) and [Twitter](#) and subscribe to our [Telegram Channel](#). Feel free to repost and share widely Global Research articles.

[Give Truth a Chance. Secure Your Access to Unchained News, Donate to Global Research.](#)

Adenomyosis is a benign estrogen-dependent uterine disorder that may cause abnormal uterine bleeding, pelvic pain and infertility

It's estimated that about 20% of women, primarily those of reproductive age, may suffer from the condition, yet there's surprisingly little awareness surrounding it

Estrogen is believed to promote the growth of adenomyosis; many people are exposed to excessive estrogen in the form of birth control, estrogen replacement therapy and even exposure to plastics

Estrogen is also carcinogenic and antimetabolic, radically reducing the ability of your mitochondria to create cellular energy

One of the most important strategies for adenomyosis — aside from avoiding estrogen and xenoestrogens — is to take natural progesterone, which is anti-estrogenic

*

Adenomyosis is a benign estrogen-dependent uterine disorder,¹ which involves symptoms such as abnormal uterine bleeding, pelvic pain and infertility.² In cases of adenomyosis, endometrial tissue, which normally lines the uterus, is found in the myometrium, or the muscular wall of the uterus.

In other words, the uterine lining grows into the uterus' muscular wall,³ causing symptoms

ranging from mild to severe. For some women, adenomyosis causes no symptoms at all while others experience debilitating pain.

It's estimated that about 20% of women, primarily those of reproductive age, may suffer from the condition,⁴ yet there's surprisingly little awareness surrounding it among health care professionals and the public alike.⁵

What Are the Signs and Symptoms of Adenomyosis?

Adenomyosis can be categorized into two main types: focal and diffuse. In focal adenomyosis, the endometrial tissue is localized to a specific area or areas within the uterine muscle. These localized regions are sometimes referred to as adenomyomas, which are similar to fibroids but contain endometrial tissue.

Diffuse adenomyosis involves a more widespread infiltration of endometrial tissue throughout the uterine muscle. This type is characterized by a uniformly thickened uterine wall. In either type, the lining of the uterus grows into the uterine wall, leading to thickening and enlargement of the uterus, which in some cases may double or triple in size.⁶

"The displaced tissue continues to act normally — thickening, breaking down and bleeding — during each menstrual cycle," Mayo Clinic notes.⁷ Common symptoms of adenomyosis include heavy menstrual bleeding, severe menstrual cramps, pelvic pain and bloating.

Some women with adenomyosis may also experience pain during intercourse. Infertility is also possible and women with adenomyosis who become pregnant are more likely to have complications such as miscarriage, pre-term delivery, preeclampsia and post-delivery bleeding.⁸

The condition can sometimes be asymptomatic, however, making diagnosis challenging. While most women who develop adenomyosis are between the ages of 40 and 50, it can develop in younger women as well. The condition is becoming more common in women in their 30s,⁹ and even teenagers can be affected. Other factors that increase the risk of adenomyosis include:

- Carrying two or more pregnancies to term¹⁰
- Prior uterine surgeries, including uterine fibroid removal, cesarean section or dilatation and curettage (D&C)
- Endometriosis

Top Causes of Adenomyosis

A combination of hormones, genetics and inflammation or injury to the area likely influence adenomyosis development. Science Alert reported:¹¹

"It is thought that the region between the endometrium and myometrium becomes damaged, either by the natural processes of the menstrual cycle, pregnancies, and childbirth, or medical procedures. In some women, damage to the endometrial tissue layer does not heal as it should, and the endometrium-like cells enter and grow

abnormally into the myometrium.”

It's possible that uterine incisions during surgery may allow for the direct invasion of endometrial cells into the uterine wall, for instance. It's also been suggested that the misplaced endometrial tissue may be deposited in the uterine wall in the fetus during development.¹²

Inflammation of the uterine lining after childbirth is another potential factor that could degrade the boundary of cells lining the uterus. Another theory suggests bone marrow stem cells in uterine muscle could trigger the condition.¹³

Estrogen, a hormone involved in the menstrual cycle, is also believed to promote the growth of adenomyosis. Therefore, conditions that result in higher levels of estrogen may be linked to an increased risk. “Regardless of how adenomyosis develops, its growth depends on the body's circulating estrogen,” Mayo Clinic explains.¹⁴

Estrogen Promotes Adenomyosis

“Estrogens have been proven to be the crucial hormones driving the growth of adenomyosis,” as noted in the *Journal of Clinical Medicine*.¹⁵ Yet, estrogen's exact role in worsening adenomyosis is still not fully understood.

In one study, researchers found that a molecule called miR-145 is much higher in adenomyosis tissue compared to healthy tissue.¹⁶ High levels of miR-145 turn on genes that cause inflammation and cell adhesion.

One important gene affected by miR-145 is CITED2, which plays a key role in stopping cell movement. The study showed that estrogen, through a specific receptor, increases miR-145 levels, helping adenomyosis to progress. Yale Medicine also explains:¹⁷

“Evidence suggests that prolonged exposure to estrogen may be a contributing factor [to uterine adenomyosis]. Women may be exposed to more estrogen if their menstrual cycles are shorter than average or if they have been pregnant two or more times.”

However, many people are exposed to excessive estrogen in the form of [birth control](#), estrogen replacement therapy and even exposure to plastics.

The Hazards of High Estrogen

While estrogen is routinely referred to as “the female hormone,” this is misleading, as it's not exclusive to women. Moreover, there's not just one estrogen but several. One of the properties of estrogens is their ability to increase cells' ability to hold water, which is why women with estrogen dominance are prone to edema (water retention). Cellular swelling is both a characteristic of the cellular stress response and a signal for cellular proliferation.

During the follicular phase of the menstrual cycle, estrogen stimulates the uterine lining and follicles to swell and multiply in preparation for the fertilization of an egg. Similarly, during and after pregnancy, breast tissue swells and grows larger to facilitate milk production. But cellular swelling and proliferation is also a hallmark of cancer. Indeed, the word oncology

comes from the Greek word “oncos,” which means swelling.

In his 1997 book, “From PMS to Menopause: Female Hormones in Context,”¹⁸ biologist Ray Peat¹⁹ stated that estrogen had been shown to replicate the shock phase of the stress reaction in animals. According to Peat, the physiological purpose of estrogens is to stimulate cell division by triggering water uptake by the cell. Peat also suspected that estrogen was a metabolic inhibitor that slows down energy production in the cell.

Further, as bioenergetic researcher Georgi Dinkov explained in my interview with him, [estrogen is carcinogenic and antimetabolic](#), radically reducing the ability of your mitochondria to create cellular energy in a form of ATP by depending on aerobic glycolysis (the Warburg effect), which radically impairs oxidative phosphorylation. This further contributes to its carcinogenic effect.

According to Dinkov, the biochemical role of estrogen is to aid in wound healing. In cases of tissue trauma, estrogen reverts the differentiated cells in that specific tissue back to a stem cell-like condition, to repair the damaged tissue. In young, healthy women, progesterone will turn off estrogen’s activity. However, progesterone declines with age, but estrogen synthesis typically does not. Hence, if your estrogen is high and progesterone low, your cancer risk will rise.

Estrogen also promotes the differentiation and proliferation of fat cells (adipocytes) in the subcutaneous fat layer, which is particularly prevalent in the thighs, hips and buttocks and therefore leads to increased fat storage in these areas.

Some people and clinicians believe that bioidentical estrogen solves the problem but it does nothing of the sort. Bioidentical estrogen still has all the same negative characteristics. It increases the risk of all cancers, lowers your metabolic rate and increases your risk of obesity, diabetes and osteoporosis — along with adenomyosis.

Tips to Lower Your Estrogen Exposure and Load

Conventional adenomyosis treatment typically involves pain medications and hormonal treatments, such as birth control pills, and even surgery, including hysterectomy. But one of the most important strategies — aside from avoiding estrogen and xenoestrogens — is to take natural progesterone. Not only is progesterone anti-estrogenic but it’s a potent cortisol blocker and will improve mitochondria production of cellular energy by blocking estrogen and cortisol.

Before you consider using progesterone it is important to understand that it is not a magic bullet and you get the most benefit by implementing a Bioenergetic diet approach that allows you to effectively burn glucose as your primary fuel with backing up electrons in your mitochondria that reduces your energy production. My new book coming out shortly about Cellular Health covers this process in great detail.

Once you have dialed in your diet, an effective strategy that can help counteract estrogen excess is to take transmucosal progesterone (not oral or transdermal), which is a natural estrogen antagonist. Progesterone is one of only four hormones I believe many adults can benefit from. (The other three are thyroid hormone T3, DHEA and pregnenolone.)

I do not recommend transdermal progesterone, as your skin expresses high levels of 5-alpha

reductase enzyme, which causes a significant portion of the progesterone you're taking to be irreversibly converted primarily into allopregnanolone and cannot be converted back into progesterone.

As a general recommendation, I recommend taking 25 to 50 mg of bioidentical progesterone per day, taken in the evening one hour before bed, as it can also promote sleep. For optimal bioavailability, progesterone needs to be mixed into natural vitamin E. The difference in bioavailability between taking progesterone orally without vitamin E and taking it with vitamin E is 45 minutes versus 48 hours.

You can make your own by dissolving pure USP progesterone powder into one capsule of a high-quality vitamin E, and then rub the mixture on your gums. Fifty milligrams of powdered progesterone is about 1/32 teaspoon.

Do not use synthetic vitamin E (alpha tocopherol acetate — the acetate indicates that it's synthetic). Natural vitamin E will be labeled "d alpha tocopherol." This is the pure D isomer, which is what your body can use.

There are also other vitamin E isomers, and you want the complete spectrum of tocopherols and tocotrienols, specifically the beta, gamma, and delta types, in the effective D isomer. As an example of an ideal vitamin E you can look at the label on our vitamin E in our store. You can use any brand that has a similar label.

If you are a menstruating woman, you should take the progesterone during the luteal phase or the last half of your cycle which can be determined by starting 10 days after the first day of your period and stopping the progesterone when your period starts.

If you are a male or non-menstruating woman you can take the progesterone every day for 4-6 months and then cycle off for one week. The best time of day to take progesterone is 30 minutes before bed as it has an anti-cortisol function and will increase GABA levels for a good night's sleep.

Please note that when progesterone is used transmucosally on your gums as I advise, the FDA believes that somehow converts it into a drug and prohibits any company from advising that on its label. However, please understand that it is perfectly legal for any physician to prescribe an off-label indication for a drug.

In this case progesterone is a natural hormone and not a drug and is very safe even at high doses. This is unlike synthetic progesterone, called progestins, that are used by drug companies, but frequently, and incorrectly, referred to as progesterone, which are dangerous and should never be used by anyone.

Considering the health risks associated with estrogen excess, here are some additional commonsense strategies that can help you limit your exposure and lower your estrogen load:

- Avoid synthetic estrogens — Minimize exposure to synthetic estrogens, such as those found in hormone replacement therapy and oral contraceptives. Consult with a qualified health care professional about alternative treatments and/or contraceptive methods with lower estrogen content.
- Avoid linoleic acid (LA) — Omega-6 PUFA like LA functions very similarly to estrogen as they both increase your risk for cancer and decrease metabolic

function by suppressing your thyroid. Best to read my [comprehensive LA article](#) for more details.

- Choose natural products — Opt for natural and organic personal care products, including makeup, skin care, and hair care items, to reduce exposure to synthetic chemicals like parabens and phthalates, which have estrogenic properties.
- Limit pesticide exposure — Choose organic produce whenever possible to reduce exposure to pesticides, many of which have estrogenic effects. Washing fruits and vegetables thoroughly can also help remove pesticide residues.
- Rethink your household products — Many household cleaning products, laundry detergents and air fresheners contain chemicals with estrogenic properties. Swap them out for natural, nontoxic alternatives or make your own cleaning solutions using vinegar, baking soda and essential oils instead.
- Avoid plastic containers — Minimize the use of plastic containers and food packaging, which can leach estrogenic compounds into food and beverages. Instead, opt for glass or stainless steel containers for food storage and water bottles.
- Maintain a healthy weight — Aim for a healthy weight and body composition through a balanced diet and regular exercise. Excess body fat, particularly around the thighs, hips, and buttocks, can contribute to higher estrogen levels.
- Support liver health — Support liver function, as your liver plays a crucial role in metabolizing and eliminating excess estrogen from your body. Eat a nutrient-rich diet, limit alcohol consumption and consider incorporating liver-supporting herbs and supplements, such as milk thistle or dandelion root.
- Reduce stress — Manage stress through relaxation techniques like meditation, deep breathing exercises, yoga or spending time in nature. Chronic stress can disrupt hormone balance, including estrogen levels, so prioritizing stress reduction is essential.

*

Note to readers: Please click the share button above. Follow us on Instagram and Twitter and subscribe to our Telegram Channel. Feel free to repost and share widely Global Research articles.

Notes

¹ [Molecular Medicine Reports March 8, 2021](#)

² [J Clin Med. 2023 Jul; 12\(14\): 4828](#)

^{3, 10, 17} [Yale Medicine, Uterine Adenomyosis](#)

⁴ [Best Practice & Research Clinical Obstetrics & Gynaecology August 2006, Volume 20, Issue 4, Pages 465-477](#)

^{5, 8, 11} [Science Alert May 13, 2024](#)

^{6, 9} [Cleveland Clinic, Adenomyosis](#)

^{7, 12, 13, 14} [Mayo Clinic, Adenomyosis](#)

¹⁵ [J Clin Med. 2022 Aug; 11\(15\): 4407](#)

¹⁶ [Reproductive BioMedicine Online May 12, 2024, 104108](#)

¹⁸ [From PMS to Menopause: Female Hormones in Context \(PDF\)](#)

¹⁹ [Ray Peat Articles](#)

Featured image is from Mercola

The original source of this article is [Mercola](#)
Copyright © [Dr. Joseph Mercola](#), [Mercola](#), 2024

[Comment on Global Research Articles on our Facebook page](#)

[Become a Member of Global Research](#)

Articles by: [Dr. Joseph Mercola](#)

Disclaimer: The contents of this article are of sole responsibility of the author(s). The Centre for Research on Globalization will not be responsible for any inaccurate or incorrect statement in this article. The Centre of Research on Globalization grants permission to cross-post Global Research articles on community internet sites as long the source and copyright are acknowledged together with a hyperlink to the original Global Research article. For publication of Global Research articles in print or other forms including commercial internet sites, contact: publications@globalresearch.ca
www.globalresearch.ca contains copyrighted material the use of which has not always been specifically authorized by the copyright owner. We are making such material available to our readers under the provisions of "fair use" in an effort to advance a better understanding of political, economic and social issues. The material on this site is distributed without profit to those who have expressed a prior interest in receiving it for research and educational purposes. If you wish to use copyrighted material for purposes other than "fair use" you must request permission from the copyright owner.

For media inquiries: publications@globalresearch.ca