

Are COVID Vaccines Causing Liver Failure?

By [Dr. Joseph Mercola](#)

Global Research, April 26, 2022

[Mercola](#)

Region: [Europe](#), [USA](#)

Theme: [Science and Medicine](#)

All Global Research articles can be read in 51 languages by activating the “Translate Website” drop down menu on the top banner of our home page (Desktop version).

To receive Global Research’s Daily Newsletter (selected articles), [click here](#).

Visit and follow us on [Instagram](#), [Twitter](#) and [Facebook](#). Feel free to repost and share widely Global Research articles.

As of April 8, 2022, 74 cases of severe hepatitis that health officials can’t explain have been reported in children up to 10 years old

In October 2021, a case report involving a 47-year-old, previously healthy, man demonstrated conclusive evidence that COVID-19 shots may trigger hepatitis

A Journal of Hepatology paper noted that seven additional cases of suspected immune-mediated hepatitis have been reported following COVID-19 shots

Researchers have uncovered innate immune suppression triggered by COVID-19 shots and other disturbances that could cause liver disease

Recently released Pfizer documents also show that, in the first week after the shot, people of all ages experienced a temporary weakening of the immune system; could this increased susceptibility to infection also be playing a role in hepatitis and other cases of liver disease?

*

A strange outbreak of severe hepatitis in young children has been reported in the U.S. and Europe, puzzling public health officials. The children were tested for common hepatitis viruses, but they were not to blame, leaving the cause unknown.

In a news release, Graham Cooke, a professor of infectious diseases at Imperial College London, suggested that if the hepatitis was caused by COVID-19, “it would be surprising not to see it more widely distributed across the country given the high prevalence of (COVID-19) at the moment.”¹

Potential links to COVID-19 injections appear not to have been widely explored yet, even though the shots have been previously associated with the development of hepatitis.² British health officials did state, however, that none of the affected children had received a

COVID-19 shot.³

Young Children Developing Mysterious Liver Disease

In the U.S., nine children in Alabama have developed severe hepatitis, or inflammation of the liver, that health officials can't explain. All of the children were ages 6 years and younger and were previously healthy.⁴ Symptoms of the liver disease include diarrhea, nausea and vomiting, along with jaundice in some. Liver enzymes were also elevated.

Five of the children tested positive for adenovirus type 41, which are respiratory viruses that can cause the common cold. Health officials have suggested that adenovirus type 41 could be to blame, but Dr. Wes Stubblefield, district medical officer for the Alabama Department of Public Health, admitted to NBC News, "This is unusual. This virus hasn't, in the past, been associated with this constellation of signs, symptoms and injury."⁵

Others have also discounted this theory, as adenoviruses are extremely common in children, meaning that it's quite possible they could test positive for adenoviruses without them being the cause of the hepatitis.⁶ As of April 8, 2022, 74 cases of hepatitis had been reported in children up to 10 years old. Some of the children required hospitalization and six have undergone liver transplants, but no deaths were reported as of April 11, 2022.

With the increase in cases reported over the last month, the World Health Organization expects that more cases of the mysterious hepatitis illness will be uncovered in children. So far, laboratory testing has ruled out hepatitis type A, B, C, and E viruses, along with hepatitis D where applicable.

"Overall," WHO reported, "the etiology of the current hepatitis cases is still considered unknown and remains under active investigation. Laboratory testing for additional infections, chemicals and toxins is underway for the identified cases."⁷

COVID-19 Shots May Trigger Hepatitis

A case report involving a 47-year-old, previously healthy man demonstrates conclusive evidence that COVID-19 shots may trigger hepatitis. "Immune-mediated hepatitis with the Moderna vaccine," researchers wrote in the *Journal of Hepatology* in October 2021, "[is] no longer a coincidence but confirmed."⁸

The man featured in the case report received his first Moderna COVID-19 shot on April 26, 2021. Three days later, he developed malaise and jaundice, a yellowing of the skin that can occur if your liver isn't processing red blood cells efficiently; it's a hallmark of hepatitis, and a symptom being experienced by some of the children noted above.

The man had his liver function tested four years earlier, with tests coming back normal, and had no history of acetaminophen usage, which can cause liver damage, and only minimal alcohol usage. Yet, three days after the shot, liver tests showed concerning results:⁹

"Investigations on the 30th April showed serum bilirubin 190 $\mu\text{mol/L}$ (normal 0-20), alanine aminotransferase (ALT) 1,048 U/L (normal 10-49), alkaline phosphatase (ALP) 229 U/L (normal 30-130) ..."

By the end of June, the man's jaundice and liver function tests improved, but then he received a second dose of the Moderna shot on July 6, 2021. Within days, the jaundice returned and liver function tests declined. "The pattern of injury on histology was consistent with acute hepatitis, with features of autoimmune hepatitis or possible drug-induced liver injury (DILI), triggering an autoimmune-like hepatitis," the researchers explained, adding:¹⁰

"This case illustrates immune-mediated hepatitis secondary to the Moderna vaccine, which on inadvertent re-exposure led to worsening liver injury with deranged synthetic function. This occurred in a well man with no other medical problems. The onset of jaundice associated with the mRNA vaccine was unusually rapid."

Hepatitis Cases Reported Following Shots

The case report above isn't an isolated one. The Journal of Hepatology paper noted that seven additional cases of suspected immune-mediated hepatitis have been reported following COVID-19 shots, including both Pfizer's and Moderna's.

They hope to raise awareness so that vaccination centers will routinely check for signs of immune-mediated hepatitis prior to administering second doses and state, "Long-term follow up of identified individuals will be essential in determining the prognosis of this immune-mediated liver injury."¹¹

In a separate letter to the editor, published in the Journal of Hepatology in June 2021, researchers again raised concerns that COVID-19 shots could cause hepatitis. In this case, a 56-year-old woman developed severe autoimmune hepatitis following her first dose of Moderna's COVID-19 shot.¹²

Prior to this, in April 2021, researchers also described a case of autoimmune hepatitis that developed after a COVID-19 shot, this time in a 35-year-old woman who was three months postpartum. In autoimmune hepatitis, the body's immune system mistakenly attacks the liver, causing inflammation and damage, and it's possible the shot triggered the autoimmunity via spike-directed antibodies:¹³

"To our knowledge, this is the first reported episode of autoimmune hepatitis developing post-COVID-19 vaccination, raising concern regarding the possibility of vaccine-induced autoimmunity. As causality cannot be proven, it is possible that this association is just coincidental.

However, severe cases of SARS-CoV-2 infection are characterized by an autoinflammatory dysregulation that contributes to tissue damage. As the viral spike protein appears to be responsible for this, it is plausible that spike-directed antibodies induced by vaccination may also trigger autoimmune conditions in predisposed individuals."

Is Immune Suppression to Blame?

Researchers from Ireland noted in November 2021, "It is speculated that SARS-CoV-2 can disturb self-tolerance and trigger autoimmune responses through cross-reactivity with host cells and that the COVID-19 mRNA vaccines may trigger the same response." They also

reported the cause of autoimmune hepatitis that developed after a COVID-19 injection in a 71-year-old woman with no risk factors for autoimmune disease.

She noticed jaundice four days after the shot and had “markedly abnormal” liver function tests. The researchers raised the possibility that this is a case of vaccine-related drug-induced liver injury and, like the other teams that reported similar cases, noted:¹⁴

“These findings raise the question as to whether COVID-19 mRNA vaccination can, through activation of the innate immune system and subsequent non-specific activation of autoreactive lymphocytes, lead to the development of autoimmune diseases including AIH or trigger a drug-induced liver injury with features of AIH.

The trigger, if any, may become more apparent over time, especially following withdrawal of immunosuppression. As with other autoimmune diseases associated with vaccines the causality or casualty factor will prove difficult to tease apart ... But it does beg the question of whether or not these individuals should receive the second dose of an mRNA COVID-19 vaccine.”

Stephanie Seneff, a senior research scientist at the Massachusetts Institute of Technology (MIT), and colleagues have also highlighted the innate immune suppression triggered by COVID-19 shots.¹⁵

mRNA COVID-19 shots teach your cells to produce a protein, or piece of protein, that triggers an immune response, including the production of antibodies.¹⁶ However, because natural mRNA is easily broken down, this means the experimental gene therapy needs a special delivery system to make it to the body’s cells.

The shots use lipid nanoparticles that contain polyethylene glycol (PEG)¹⁷ for this purpose. The mRNA is wrapped in lipid nanoparticles (LNPs) that carry it to your cells, and the LNPs are “PEGylated” — that is, chemically attached to PEG molecules to increase stability.¹⁸

Usually, if you were to inject RNA into your body, enzymes would immediately break it apart, but the COVID-19 shots are specifically designed so that doesn’t happen. As such, “mRNA vaccines promote sustained synthesis of the SARS-CoV-2 spike protein,” Seneff and colleagues write in *Food and Chemical Toxicology*.¹⁹ The spike protein is not only neurotoxic but also impairs DNA repair mechanisms, while suppression of type I interferon responses result in impaired innate immunity, they explain.²⁰

COVID Shot Disturbances Could Cause Liver Disease

Seneff’s research suggests genetic modifications introduced by COVID-19 shots may induce immune cells to release large quantities of exosomes into circulation. Exosomes are extracellular vesicles that contain protein, DNA, RNA and other constituents, and may contain mRNA along with spike protein. According to Seneff and colleagues:²¹

“[W]e present evidence that vaccination induces a profound impairment in type I interferon signaling, which has diverse adverse consequences to human health. Immune cells that have taken up the vaccine nanoparticles release into circulation large numbers of exosomes containing spike protein along with critical microRNAs that

induce a signaling response in recipient cells at distant sites.

We also identify potential profound disturbances in regulatory control of protein synthesis and cancer surveillance. These disturbances potentially have a causal link to neurodegenerative disease, myocarditis, immune thrombocytopenia, Bell's palsy, liver disease, impaired adaptive immunity, impaired DNA damage response and tumorigenesis."

In one example noted in their study, the shot appears to have caused a case of viral reactivation that led to liver failure. The case involved an 82-year-old woman who had hepatitis C (HCV) in 2007. Just days after she received a Pfizer COVID-19 shot, "a strong increase in HCV load occurred," along with jaundice. She died from liver failure three weeks after the injection.²²

They also report that the release of exosomes containing microRNAs following COVID-19 shots could interfere with IRF9 synthesis, leading to reduced synthesis of sulfatide in the liver. This cascade, they believe, could represent a "plausible factor" in the multiple case reports that have found liver damage following COVID-19 shots.²³

When they reviewed data from the Vaccine Adverse Event Reporting System (VAERS), including symptoms that "clearly represent serious liver problems," they identified 731 events following COVID-19 shots — representing more than 97% of cases out of all vaccines in 2021.²⁴

Pfizer documents released by the U.S. Food and Drug Administration in April 2022 must also be taken into account. Buried in one of the documents is the statement, "Clinical laboratory evaluation showed a transient decrease in lymphocytes that was observed in all age and dose groups after Dose 1, which resolved within approximately one week ..."²⁵

What this means is Pfizer knew that, in the first week after the shot, people of all ages experienced transient immunosuppression, or put another way, a temporary weakening of the immune system, after the first dose. Could this increased susceptibility to infection also be playing a role in hepatitis and other cases of liver disease following the shots? An investigation is warranted to find out.

*

Note to readers: Please click the share buttons above or below. Follow us on Instagram, Twitter and Facebook. Feel free to repost and share widely Global Research articles.

Notes

^{1, 3, 6} [Yahoo April 19, 2022](#)

^{2, 8} [Journal of Hepatology October 4, 2021](#)

^{4, 5} [NBC News April 15, 2022](#)

⁷ [WHO April 15, 2022](#)

⁹ [Journal of Hepatology October 4, 2021, Case description](#)

¹⁰ [Journal of Hepatology October 4, 2021, Case description, Discussion](#)

¹¹ [Journal of Hepatology October 4, 2021, Discussion](#)

¹² [Journal of Hepatology June 17, 2021](#)

¹³ [Journal of Hepatology April 13, 2021](#)

¹⁴ [J Hepatol. 2021 Nov; 75\(5\): 1252-1254](#)

^{15, 19, 21} [Food and Chemical Toxicology June 2022, Volume 164, 113008](#)

¹⁶ [U.S. CDC December 18, 2020](#)

¹⁷ [Anthrax Vaccine January 11, 2021](#)

¹⁸ [Science December 21, 2020](#)

²⁰ [Food and Chemical Toxicology June 2022, Volume 164, 113008, Highlights](#)

²² [Food and Chemical Toxicology June 2022, Volume 164, 113008, Section 9](#)

²³ [Food and Chemical Toxicology June 2022, Volume 164, 113008, Section 11](#)

²⁴ [Food and Chemical Toxicology June 2022, Volume 164, 113008, Section 15.2](#)

²⁵ [The Naked Emperor Substack March 29, 2022](#)

Featured image is from The Conservative Woman

The original source of this article is [Mercola](#)
Copyright © [Dr. Joseph Mercola](#), [Mercola](#), 2022

[Comment on Global Research Articles on our Facebook page](#)

[Become a Member of Global Research](#)

Articles by: **[Dr. Joseph Mercola](#)**

Disclaimer: The contents of this article are of sole responsibility of the author(s). The Centre for Research on Globalization will

not be responsible for any inaccurate or incorrect statement in this article. The Centre of Research on Globalization grants permission to cross-post Global Research articles on community internet sites as long the source and copyright are acknowledged together with a hyperlink to the original Global Research article. For publication of Global Research articles in print or other forms including commercial internet sites, contact: publications@globalresearch.ca

www.globalresearch.ca contains copyrighted material the use of which has not always been specifically authorized by the copyright owner. We are making such material available to our readers under the provisions of "fair use" in an effort to advance a better understanding of political, economic and social issues. The material on this site is distributed without profit to those who have expressed a prior interest in receiving it for research and educational purposes. If you wish to use copyrighted material for purposes other than "fair use" you must request permission from the copyright owner.

For media inquiries: publications@globalresearch.ca