

CDC Data Shows Links Between Vaccines and Sudden Infant Death Syndrome (SIDS)

749 deaths linked to Hib (Haemophilus influenzae Type b) and 51% were SIDS

By [Sayer Ji](#) and [Dr. William Makis](#)

Global Research, March 27, 2024

[COVID Intel and GreenMedInfo](#)

Region: [USA](#)

Theme: [Science and Medicine](#)

All Global Research articles can be read in 51 languages by activating the Translate Website button below the author's name (only available in desktop version).

To receive Global Research's Daily Newsletter (selected articles), [click here](#).

Click the share button above to email/forward this article to your friends and colleagues. Follow us on [Instagram](#) and [Twitter](#) and subscribe to our [Telegram Channel](#). Feel free to repost and share widely Global Research articles.

[Global Research Fundraising: Stop the Pentagon's Ides of March](#)

By [Sayer Ji](#), [GreenMedInfo](#)

The CDC's own research found that the long denied vaccine-SIDS link is real.

If you believe the official pronouncements of top governmental health agencies like the CDC and FDA, all the vaccines in the present day schedule are a *priori* safe and effective.

Not only are you told that they can't harm you, but that not taking them can kill you.

Parents are under even more pressure. **They are told that refraining from vaccinating their infants or children will greatly increase their risk of dying or being disabled.** Worse, they are increasingly labeled as 'crazy' and 'irresponsible' anti-vaccine zealots who are putting the lives of others in danger.

But what happens when the actual evidence from the scientific and clinical literature produced by these very agencies contradicts their own vaccine policies?

This is exactly what has happened with the publication of a new study in the Journal of Pediatrics titled ,

["Adverse Events following Haemophilus influenzae Type b Vaccines in the Vaccine Adverse Event Reporting System, 1990-2013,"](#) wherein **CDC and FDA researchers identify 749 deaths linked to the administration of the Hib vaccine, 51% of which were sudden infant death linked to the administration of Hib vaccine.**

The CDC has boldly denied that there is any evidence supporting a causal link between vaccines and infant death, despite the fact that their own webpage on the topic acknowledges that **“From 2 to 4 months old, babies begin their primary course of vaccinations. This is also the peak age for sudden infant death syndrome (SIDS).”** **Written off as coincidence, the CDC suggests that stomach sleeping is the primary modifiable risk factor.**

Because [SIDS is the 3rd leading cause of death in infants](#), and because the U.S. has one of the [highest infant mortality rates in the developed world](#), one would think that more progress would have been made toward understanding its causes. Perhaps, as [explored in this past article](#), the signal of harm is being ignored. Neglect and suppression of available data has recently been exposed with the confession of a top CDC vaccine scientist who was compelled to covered up data revealing an [autism-MMR link in African-American boys](#).

In the new study, the CDC and FDA researchers themselves acknowledge **“the scarcity” of post licensure safety data on Hib vaccines in today’s vaccination schedule.** They evaluated reports involving the currently licensed Hib vaccines received from January 1, 1990, through December 1, 2013 available on the Vaccine Adverse Event Reporting System (VAERS).

Presently, the CDC recommends 4 doses of the Hib vaccine at the following ages: 2 months, 4 months, 6 months, 12 months through 15 months.

The Hib vaccine is [described on the CDC website](#) as “very safe” and “effective” at preventing Hib disease, which it states can be deadly. They list “most common side effects as usually mild and last 2 or 3 days,” including “redness, swelling, and warmth where the child got the shot” and “fever”. Nowhere is there listed death or disability as a possible side effect.

In stark contrast to these statements the study uncovered the following highly concerning results:

VAERS received 29,747 reports after Hib vaccines; 5179 (17%) were serious, including 896 reports of deaths. Median age was 6 months (range 0-10.22 months). Sudden infant death syndrome was the stated cause of death in 384 (51%) of 749 death reports with autopsy/death certificate records. The most common non-death serious AE categories were **neurologic** (80; 37%), other noninfectious (46; 22%) (comprising mainly constitutional signs and symptoms); and gastrointestinal (39; 18%) conditions. No new safety concerns were identified after clinical review of reports of AEs that exceeded the data mining statistical threshold.

Consider also that VAERS is a passive surveillance system, which suffers from profound underreporting. According to the VAERS [site’s own disclaimer](#):

“Underreporting” is one of the main limitations of passive surveillance systems, including VAERS. The term, underreporting refers to the fact that VAERS receives reports for only a small fraction of actual adverse events. The degree of underreporting varies widely. As an example, a great many of the millions of vaccinations administered each year by injection cause soreness, but relatively few of these episodes lead to a VAERS report.

According to Barbara Loe Fisher, founder of the [National Vaccination Information Center](#), underreporting may result in overlooking 99% or higher of all vaccine associated injuries:

Former FDA Commissioner David Kessler estimated in a 1993 article in the Journal of the American Medical Association that **fewer than 1 percent of all doctors report injuries and deaths following the administration of prescription drugs**. This estimate may be even lower for vaccines. In one [survey](#) that our organization conducted in New York in 1994, only 1 doctor in 40 reported to VAERS.

Considering the influence of underreporting, these deaths represent only the tip of the iceberg of **vaccine-induced infant morbidity and mortality caused by Hib vaccines**. The study also mentioned an earlier analysis which found that infant death is the most common cause of death reported by all vaccine linked reports on VAERS, **“accounting for almost one-half of all deaths reported.”**

Obviously, this is an appalling study. The death of even 1 child for a potentially ineffective medical intervention designed to prevent a [rarely fatal](#) illness is a tragedy. Nor can any single vaccine be proven to have prevented any single case of disease because the clinical outcome (end point) is a non-event. This is not the case, however, for vaccine side effects which can be linked directly to the vaccination event with plausible scientific mechanisms.

What is perhaps most astounding is the researcher’s conclusion:

Review of VAERS reports did not identify any new or unexpected safety concerns for Hib vaccines.

This callous disregard for the evidence — **evidence that clearly shows the CDC misrepresents the safety of the Hib vaccine** — speaks to the blind investment in vaccine policy decisions over human wellbeing. Millions of parents have listened to the CDC and FDA and believed that these vaccines not only work but are safe. Informed consent requires those undergoing a quasi-mandatory medical intervention like vaccination to know the true risks associated with it. Failing to do so is clearly a violation of this medical ethical protection against being abused, and in some cases disabled and even killed.

My Take...

VAERS has 1881 deaths of babies 0 to 2 years old after Hib Vaccine.

1514 deaths are < 6 months.

Search Results

From the 2/23/2024 release of VAERS data:

Found 1,881 cases where Age is under-2 and Vaccine is HBHEPB or HBPV or HIBV and Patient Died

Government Disclaimer on use of this data

Table

↓	↑ ↓	Count	Percent
Age			
< 6 Months		1,514	80.49%
6-11 Months		211	11.22%
1-2 Years		156	8.29%
TOTAL		1,881	100%

Case Details

This is page 1 out of 189

*

Note to readers: Please click the share button above. Follow us on Instagram and Twitter and subscribe to our Telegram Channel. Feel free to repost and share widely Global Research articles.

Dr. William Makis is a Canadian physician with expertise in Radiology, Oncology and Immunology. Governor General’s Medal, University of Toronto Scholar. Author of 100+ peer-reviewed medical publications.

Featured image is from COVID Intel

The original source of this article is [COVID Intel and GreenMedInfo](#)
Copyright © [Sayer Ji](#) and [Dr. William Makis](#), [COVID Intel and GreenMedInfo](#), 2024

[Comment on Global Research Articles on our Facebook page](#)

[Become a Member of Global Research](#)

Articles by: [Sayer Ji](#) and [Dr. William Makis](#)

Disclaimer: The contents of this article are of sole responsibility of the author(s). The Centre for Research on Globalization will not be responsible for any inaccurate or incorrect statement in this article. The Centre of Research on Globalization grants permission to cross-post Global Research articles on community internet sites as long the source and copyright are acknowledged together with a hyperlink to the original Global Research article. For publication of Global Research articles in print or other forms including commercial internet sites, contact: publications@globalresearch.ca
www.globalresearch.ca contains copyrighted material the use of which has not always been specifically authorized by the copyright owner. We are making such material available to our readers under the provisions of "fair use" in an effort to advance a better understanding of political, economic and social issues. The material on this site is distributed without profit to those who have expressed a prior interest in receiving it for research and educational purposes. If you wish to use copyrighted material for purposes other than "fair use" you must request permission from the copyright owner.

For media inquiries: publications@globalresearch.ca