

# COVID-19 Vaccine Induced Psychosis - 13 Cases of Post-Vaccine Psychosis, Mania & Suicide Attempts That Will Shock You.

By Dr. William Makis

Global Research, March 09, 2023

**COVID Intel 8 March 2023** 

Theme: Science and Medicine

All Global Research articles can be read in 51 languages by activating the **Translate Website** button below the author's name (desktop version)

To receive Global Research's Daily Newsletter (selected articles), click here.

Follow us on <u>Instagram</u> and <u>Twitter</u> and subscribe to our <u>Telegram Channel</u>. Feel free to repost and share widely Global Research articles.

\*\*\*

COVID-19 vaccines can significantly impact mental health. Family members have reported brain fog, mood changes, personality changes, hostility and anger, erratic driving, confusion and more. Sometimes, the impact on mental health is catastrophic.

Here are 13 cases that will give you a new perspective on what COVID-19 vaccines can do to the mental health of a person with no prior psychiatric history.

#### Case 1: 31 yo Hispanic man has psychosis after Moderna

A 31-year-old, single Hispanic man without past medical or psychiatric history, was brought to emergency by police because of erratic and bizarre behavior (Reinfeld et al.)

He was found to be anxious, guarded, superficial and grandiose. He reported becoming 'clairvoyant', being able to talk with dead people, hearing 'people drumming outside his house' and the constant voice of a co-worker whom he believed to be a paramour, but was not.

All these symptoms began one month ago, after receiving the first dose of Moderna COVID-19 vaccine, and markedly worsened three weeks later after receiving the second dose. Previously, he was working full-time as an office manager.

## Case 2: 45 yo Asian-American married woman, has psychosis after Moderna

45 yo woman with no psychiatric history, presented to psychiatry after having **psychosis for 3 months**. (Alphonso et al.)

Her new-onset paranoia and auditory hallucinations began 1 month after receiving her second Moderna COVID-19 vaccine. She abruptly quit her job of 18 years and stopped eating—losing 56 lb in 4 months.

The patient would only get dressed in the dark, believing people could see her in her closet or bathroom behind a closed door. She would pace all day and stopped doing the household chores she had done for years.

She would wake her husband in the middle of the night trying to convince him that the neighbors were going to hurt her and were breaking into their home. She became angry when he tried to convince her these things were not real. Due to this irrational fear, she would sleep only 2 to 3 hours per night and no longer slept in the bedroom, preferring to lock herself in the bathroom to sleep.

She heard whispers of people talking about her and would carry on full conversations with them. She would only do this in a room alone with the door closed. Family would ask her who she was talking to, and she would become angry.

#### Case 3: 37 yo Japanese man has psychosis after Moderna

A 37 yo Japanese man with no psychiatric history had Moderna booster shot and complained of headache, a floating sensation and difficulty concentrating. (<u>Kita et al</u>)

He presented 4 days later with **talkativeness**, and grandiose delusions, saying that he had won 2 billion yen in horse racing. He also presented with emotional instability, such as crying, sleeplessness, excitement, hyperactivity and sexual deviance.

He was discharged. Nine days after Moderna booster **he jumped from the 2nd floor of his house**, and was brought back to hospital by ambulance. He exhibited flight of ideas, hyperactivity, distraction, hyperthymia and **religious delusions**, **such as saying "my child is God"**. He displayed lack of insight and became **enraged** when his actions were restricted.

He was diagnosed with acute mania with psychotic features and stayed **66 days** in the hospital.

## Case 4: 42 yo Turkish man has psychosis & mania after Pfizer

42 yo man with no psychiatric history was admitted to psychiatric emergency 5 days after 1st dose of Pfizer with complaints of **irritability and sleeplessness**. (Yesilkaya)

The patient had delusions that his family was being followed by the deep state and that they were in danger.

At clinical examination, his speech output, speed as well as psychomotor activity were increased. The patient's affect was **anxious and mood was dysphoric**. He exhibited a loosening of associations, described **persecutory and reference delusions** and displayed a lack of insight. The Young Mania Rating Scale (YMRS) score was 45.

At the routine psychiatric evaluation 15 days after discharge, the patient declared that he was unable to remember the initiation of psychiatric symptoms.

## Case 5: 45 yo Latvian man has psychosis & suicide attempt after mRNA

45 yo Latvian man with no psychiatric history presented 1 month after 2nd dose of mRNA vaccine, accompanied by his parents because of **bizarre behavior and an attempted suicide by hanging in the early morning.** (Renemane et al.)

Immediately after his 2nd mRNA dose, he developed insomnia, unreasonable anxiety and tremor.

After 2 weeks, he realized that he has been jinxed as he found some white powder under the carpet in his apartment. From that moment, he became cautious, did not leave the apartment, and reported persecution.

On the last day before his visit to the psychiatric clinic, the patient saw a man walking past the windows of his apartment and watching him. The patient described the thoughts in his head as not his own, giving him commands to observe the person on the street. He did not sleep that night and had a strong belief that he should commit suicide. He attached a rope, tried to hang himself, but his father stopped him.

In psychiatric department, he was found to be **emotionally withdrawn, immersed in his experiences, hypomimic (he sat in one position, spoke very quietly, did not actively participate in the conversation, answered questions in monosyllables or in short sentences after a long pause)**. He was disoriented to time and reported he was full of fear. He was hospitalized for 4 weeks.

#### Case 6: 20 yo woman has psychosis & seizure after Pfizer

20 yo woman presented to ER with urinary frequency, anxiety, insomnia and a fixation that she suffered from irritable bowel and kidney disease 1 week after 1st dose of Pfizer (Flannery et al.)

She displayed waxing and waning **hypochondriacal delusions that she had contracted COVID-19 and that "her body was shutting down.**" The patient was also noted to have some motor dysfunction and a transient bout of aphasia.

She also had accusatory auditory hallucinations but denied suicidal or homicidal ideation. She was kept overnight. The following morning the patient removed her clothing and had a bowel movement on the floor. She was admitted and put on medication, but became increasingly psychotic, then had a grand mal seizure and was admitted to ICU.

She also exhibited symptoms of ongoing **catatonia**, answered questions in short sentences in a **monotonous tone with low phonation**. She could ambulate, but slowly and with a one-person assist mainly to aid with initiation of movement.

She was diagnosed with **autoimmune encephalitis and spent 45 days** in the hospital.

## Case 7: 38 yo Brazilian woman has refractory psychosis after Pfizer

38 yo Brazilian woman with no psychiatric history, developed fever, shivering and myalgia

24hr after 1st Pfizer dose. Within a week she became agitated and did not recognize relatives. (Neves et al.)

She became aggressive, and presented disheveled with poor hygiene, unfriendly face, lacking self-orientation and refusing to talk to staff and uncooperative with doctors. Her thought was delusional. She believed staff were persecuting her, that several city mayors were her parents. Her mood was dysphoric and enraged.

She was put on psychiatric medications but didn't improve. After 6 months of hospitalization, she remained hostile, persecutory and refusing to talk to staff and her psychosis was defined as refractory. She never recovered.

#### Case 8: 15 yo Taiwanese boy has psychosis after Pfizer

15 yo Taiwanese boy with no psychiatric history presented to ER with **agitation**, **involuntary limb stretching and screaming** 2 days after 2nd dose of Pfizer (<u>Yen-Li Lien</u> et al.)

After admission, he was observed to demonstrate bizarre behaviors, including sitting up and lying down frequently and taking up the mannerism of praying in bed.

We also noted auditory hallucinations and delusions. After discharge, his hallucinations and bizarre behavior persisted for > 1 month, requiring additional treatment with steroids.

## Case 9: 51 yo Caucasian man, married with kids, has psychosis after AstraZeneca

The patient had his first dose of AstraZeneca COVID-19 vaccine in mid March 2021. (Roberts et al.)

Ten days later, his wife observed that he was becoming confused, with a significant change in his behavior. He was taken to the emergency department of the local general hospital by ambulance **after he stopped eating, drinking and communicating**.

He was **confused and disoriented in time and place**. His communication ranged from speaking in short sentences, sometimes **whispering or mumbling to being completely mute**. He was thought disordered. When he was given some paper to write down his thoughts, **he was unable to write anything meaningful and was drawing lines only**.

He stated that he was hearing voices, both male and female, saying the word 'Covid' to him, but he could not elaborate any further. His mood was labile and informed his behavior, which was bizarre and at times very disinhibited.

On one occasion he was incontinent of urine.

#### Case 10: 18 yo student has psychosis after AstraZeneca

18 female student with no psychiatric history was brought to emergency by family members for irrelevant talk and bizarre behavior. (Grover et al.)

She received her first dose of the vaccine, and within 2–3 h she developed high grade **fever** 

with chills, and multiple episodes of loose stools. These symptoms were followed by dizziness and one episode of fall.

As per family members fever subsided the next day, but patient reportedly started to remain anxious. She was not able sleep at night, frequently woke-up, appeared fearful and would appear to be staring in between. She would keep on pacing around and would appear distressed. Over the next 3 to 4 days, she started to remain irritable, talk irrelevantly, try to run away from home, voiced delusions of persecution and reference, and reported visual hallucinations of seeing Gods and demons.

In view of the continued symptoms and emergence of catatonia in the form of mutism, staring, rigidity and negativism, she was started on lorazepam.

#### Case 11: 22 yo woman has psychosis after AstraZeneca

22 yo woman of Middle Eastern origin presented to hospital with 3 week history of **frontal headache and fatigue** a few days after 2nd dose of AstraZeneca. (<u>Takata et al.</u>)

Symptoms did not respond to tylenol and progressed to a two-day history of confusion and hallucinations (visual and tactile). **She reported seeing "disco balls" and experiencing someone touching her skin**.

She was alert but disorientated to time, person and place, and agitated with a labile affect; she complained of auditory and visual hallucinations – for example, that the room was on fire – and delusions that were often hyper-religious in nature. She was also seen to be gesturing to the air as if responding to unseen stimuli.

After discharge from hospital, her family report that **she has not recovered** back to her pre-morbid state. **She is quieter and more withdrawn now, and still has residual psychotic symptoms of occasionally seeing green lights and praying more than before**.

## Case 12: 45 yo Croatian man has psychosis & suicide attempt after AstraZeneca

45 yo Croatian man developed unusual behavior 5 days after AstraZeneca. He became anxious, suspicious, paranoid, disorganized and complained of headaches. (<u>Borovina</u>)

Two weeks after COVID-19 vaccination, persecutory delusions and delusions of reference led him to suicide attempt by stabbing himself in the abdomen.

He had surgery and spent 23 days in the hospital.

# Case 13: 46 yo Indian teacher has psychosis after Covishield (AstraZeneca)

46 yo single mother and teacher was brought to ER by her daughter with complaints of muttering to herself, decreased sleep and appetite, hearing the voice of her husband who died 15 years back and fearful that something might happen to her and her daughter (Krishna et al.)

On repeated asking why this may happen, she explained that her husband's voice is

telling her that they would die of an accident and this would happen in the near future. Her family members noticed that she started to mutter to herself, her personal care was reduced, she stopped going to her work, and mostly preferred to stay at home for the past 20 days.

These symptoms started after 3 days of receiving the second dose of the Covishield (AstraZeneca) vaccine.

During clinical examination her affect was restricted to dull, her psychomotor activity slow, she frequently got distracted, **muttering to herself as she could hear her husband's voice talking to her during the interview** and also could not pay attention to the questions asked.

#### My Take...

These cases paint a picture that doesn't require further comment.

It is important to note, however, that COVID-19 vaccine spike protein does localize to the brain, where it can cause severe inflammation that may be autoimmune in nature (click here).

Such a tragic case of **autoimmune encephalitis after Pfizer** vaccine was reported in a 35 year old Swiss woman who developed fever, rash, behavioral changes and severe recurrent seizures 8 days after her 2nd Pfizer dose (Werner et al).

She was in and out of hospital of two months and never recovered, as she now has a seizure disorder requiring medication for life.


These episodes of psychosis have occurred after Pfizer, Moderna and AstraZeneca. In many cases, the vaccinated person was completely fine after 1st or 2nd dose but had a psychosis after 2nd dose or booster shot.

\*

Note to readers: Please click the share buttons above. Follow us on Instagram and Twitter and subscribe to our Telegram Channel. Feel free to repost and share widely Global Research articles.

**Dr. William Makis** is a Canadian physician with expertise in Radiology, Oncology and Immunology. Governor General's Medal, University of Toronto Scholar. Author of 100+ peer-reviewed medical publications.

Featured image is from the author



#### The Worldwide Corona Crisis, Global Coup d'Etat

#### **Against Humanity**

#### by Michel Chossudovsky

Michel Chossudovsky reviews in detail how this insidious project "destroys people's lives". He provides a comprehensive analysis of everything you need to know about the "pandemic" — from the medical dimensions to the economic and social repercussions, political underpinnings, and mental and psychological impacts.

"My objective as an author is to inform people worldwide and refute the official narrative which has been used as a justification to destabilize the economic and social fabric of entire countries, followed by the imposition of the "deadly" COVID-19 "vaccine". This crisis affects humanity in its entirety: almost 8 billion people. We stand in solidarity with our fellow human beings and our children worldwide. Truth is a powerful instrument."

<u>Pilots and Flight Attendants Who Recently Had Cardiac Arrests In-flight and "Died Suddenly"...</u>

ISBN: 978-0-9879389-3-0, Year: 2022, PDF Ebook, Pages: 164, 15 Chapters

Price: \$11.50 Get yours for FREE! Click here to download.

We encourage you to support the eBook project by making a donation through Global Research's **DonorBox "Worldwide Corona Crisis" Campaign Page.** 

The original source of this article is <u>COVID Intel</u> Copyright © <u>Dr. William Makis</u>, <u>COVID Intel</u>, 2023

#### **Comment on Global Research Articles on our Facebook page**

#### **Become a Member of Global Research**

Articles by: **Dr. William** 

Makis

**Disclaimer:** The contents of this article are of sole responsibility of the author(s). The Centre for Research on Globalization will not be responsible for any inaccurate or incorrect statement in this article. The Centre of Research on Globalization grants permission to cross-post Global Research articles on community internet sites as long the source and copyright are acknowledged together with a hyperlink to the original Global Research article. For publication of Global Research articles in print or other forms including commercial internet sites, contact: <a href="mailto:publications@globalresearch.ca">publications@globalresearch.ca</a>

www.globalresearch.ca contains copyrighted material the use of which has not always been specifically authorized by the copyright owner. We are making such material available to our readers under the provisions of "fair use" in an effort to advance a better understanding of political, economic and social issues. The material on this site is distributed without profit to those who have expressed a prior interest in receiving it for research and educational purposes. If you wish to use copyrighted material for purposes other than "fair use" you must request permission from the copyright owner.

For media inquiries: <a href="mailto:publications@globalresearch.ca">publications@globalresearch.ca</a>