

# Influence of Monsanto's Roundup Herbicide (Glyphosate) on Growth of Pathogenic Bacteria: Study

Influence of glyphosate in planktonic and biofilm growth of *Pseudomonas aeruginosa*

By [Global Research News](#)

Global Research, January 02, 2015

[Brazilian Journal of Microbiology](#) 9 October

2014

Theme: [Biotechnology and GMO](#),  
[Environment](#), [Science and Medicine](#)

By [Ilana Schneider Lima](#), [Nicole Carmo Baumeier](#), [Rosimeire Takaki Rosa](#), [Patrícia Maria Stuelp Campelo](#), and [Edvaldo Antonio Ribeiro Rosa](#)

## Excerpt

*Pseudomonas aeruginosa* is a gram-negative organism associated with a myriad of diseases varying from mild cutaneous infections to complicated pulmonary complications of cystic fibrosis ([Govan and Deretic, 1996](#)). It is a free-living bacterium naturally occurring in water. Surface waters can be contaminated by glyphosate due to the overuse of herbicides and to rainfalls. Due to its similarity with amino acids, it is feasible to assume that exposure to glyphosate can interfere in bacterial metabolism with shifts in biological and/or virulence characteristics.

Given the significant medical importance of *P. aeruginosa* and the fact that it is a water-borne organism, it is justifiable a study to prospect the influence of this and other herbicides, as well as other classes of pesticides, in the biology of bacteria and in the pathogenesis of their related infections.

## Abstract

This study evaluated the impact of different concentrations of glyphosate (Roundup®) on planktonic and biofilm growth of *P. aeruginosa*. Aerobic and anaerobic cultures of *P. aeruginosa* ATCC®15442 inoculated in MHB + glyphosate (0.845 ppm, 1.690 ppm, 8.45 ppm, 16.90 ppm, 84.50 ppm, 169 ppm, 845 ppm, and 1690 ppm) and cultured in normoxia and anoxia, following their OD560nm every hour for 24 h. Biofilms of adapted cells were formed in the presence of glyphosate (0.845 to 1690 ppm) in normoxia and anoxia for 36 h. Glyphosate at concentrations higher than 84.5 ppm reduces the cell density of planktonic aerobic cultures ( $p < 0.05$ ). However, these same concentrations favor the planktonic anaerobic growth ( $p < 0.05$ ). On the other hand, the herbicide favors a slight growth of biofilms in a concentration-dependent manner up to 84.5 ppm ( $p > 0.05$ ), and more pronounced over 169 ppm. Anaerobic biofilms have their growth more readily favored ( $p < 0.05$ ), regardless of concentration. In a concentration-dependent manner, glyphosate

interferes with the growth ability of *P. aeruginosa* ATCC®15442.

Keywords: *Pseudomonas aeruginosa*, glyphosate, growth, biofilm

## Introduction

Glyphosate (N-phosphonomethylglycine, CAS N° 1071-83-6) is a non-selective systemic herbicide that is widely used to control post-emergent weeds in plantations and commercial forests. Worldwide, the best known formulation is produced under the trade name Rondup®, although there are many generic products. In Brazil, the commercial formulation of Rondup® consists of 36% glyphosate isopropilaminic salt (active ingredient) and 18% polioxyethylenamin (POEA-surfactant, CAS N° 61791-26-2). Other formulations, such as Rodeo® do not have POEA in their formulation ([Tsui et al., 2004](#)).

Although the discussion about its role as a pollutant is relevant and there is a concern about possible risks of intoxication ([Stella and Ryan, 2004](#)) and genotoxicity ([Dallegrave et al., 2007](#)), very little attention has been directed to possible interactions with autochthonous human/animal microbiota, especially that with pathogenic potential. Studies to assess any influence of glyphosate or its associations on microorganisms are focused in soil and aquatic microbiomes showing a variable capacity to increase or decrease specific microbial groups ([Araújo et al., 2003](#); [Means et al., 2007](#); [Pérez et al., 2007](#)).

*Pseudomonas aeruginosa* is a gram-negative organism associated with a myriad of diseases varying from mild cutaneous infections to complicated pulmonary complications of cystic fibrosis ([Govan and Deretic, 1996](#)). It is a free-living bacterium naturally occurring in water. Surface waters can be contaminated by glyphosate due to the overuse of herbicides and to rainfalls. Due to its similarity with amino acids, it is feasible to assume that exposure to glyphosate can interfere in bacterial metabolism with shifts in biological and/or virulence characteristics.

Given the significant medical importance of *P. aeruginosa* and the fact that it is a water-borne organism, it is justifiable a study to prospect the influence of this and other herbicides, as well as other classes of pesticides, in the biology of bacteria and in the pathogenesis of their related infections.

As studies evaluating the herbicide-pathogenic microbiota interactions of are virtually nonexistent, the proposition of this study was to assess the impact that the commercial product Rodeo® exerts on *P. aeruginosa* ATCC®15442 regarding to its cell viability, adhesion ability, and biofilm formation.

## Material and Methods

### Interference on planktonic growth

*Pseudomonas aeruginosa* ATCC®15442 was grown in 1:10 Mueller-Hinton Broth plus glyphosate (at 0.845 ppm, 1.690 ppm, 8.45 ppm, 16.90 ppm, 84.50 ppm, 169 ppm, 845 ppm, and 1690 ppm) in normoxia and anoxia (90% N<sub>2</sub>, 10% CO<sub>2</sub>) until an OD<sub>560</sub> of 0.50. This step was repeated twice, to allow to cells to adapt to the distinct atmospheric conditions and to the xenobiotic presence. After adaptation, aliquots of 200 µL of suspensions containing ca. 1×10<sup>7</sup> cfu.mL<sup>-1</sup> were subcultured into tubes containing 2 mL of

1:10 Mueller-Hinton Broth plus glyphosate (at 0.845 ppm, 1.690 ppm, 8.45 ppm, 16.90 ppm, 84.50 ppm, 169 ppm, 845 ppm, and 1690 ppm). The tubes were adapted to a blood rotative mixer and incubated at 37 °C and 10 rpm, in anoxic and normoxic conditions. Growth curves were built after spectrophotometric readings at 560 nm taken every hour for twenty-four hours without interruption.

The anoxic technique used was that of [Wolfe \(2011\)](#). Anoxic conditions were maintained by displacing all air in culture broth with CO<sub>2</sub> made O<sub>2</sub> free by passage over heated copper. Mueller-Hinton Broth used for anoxic cultures was not pre-reduced. It was autoclaved and, still warm, it was aseptically vacuum-degassed under vigorous shaking. Once at room temperature, flask's headspace was filled with sterile N<sub>2</sub> and sterile glyphosate was added. Broth was carefully transferred to sterile screw capped culture tubes bearing butyl rubber seals in which bacterium inocula were previously dispensed (200 µL of suspensions containing  $ca. 1 \times 10^7$  cfu.mL<sup>-1</sup>).

To ascertain that turbidity was dependent only to cell density and not to extracellular polymeric substances, parallel anoxic/normoxic incubations were conducted with extra tubes. Each 4-h of incubation, 20 µL aliquots were harvested, streaked onto glass slides and stained with Congo Red. In all cases, optical microscopy revealed that no extracellular matter has been produced (data not shown).

#### Interference in the formation of biofilms

Aerobic or anaerobic cell suspensions of *P. aeruginosa* ATCC®15442 containing  $ca. 1 \times 10^8$  cfu.mL<sup>-1</sup> were adhered onto 96-well polystyrene wells coated with sterile 0.1% L-asparagine for 4 h at 37 °C. Polystyrene attached L-asparagine acts as anchoring points to bacterial fimbriae. Anoxic incubations were carried out in sealed jars connected to an *in house* evacuation-repletion gas system (1st and 2nd injections with N<sub>2</sub> and 3th injection with 90% N<sub>2</sub>/10% CO<sub>2</sub>).

After the period of adhesion, supernatants were removed and wells were washed with sterile 145 mM NaCl to remove non-adhered weakly adhered cells. Wells received 200 µL aliquots of 1:10 Mueller-Hinton Broth plus glyphosate (at 0.845 ppm, 1.690 ppm, 8.45 ppm, 16.90 ppm, 84.50 ppm, 169 ppm, 845 ppm, and 1690 ppm). Wells were incubated in anoxic or normoxic atmosphere for 24 h and 37 °C. Wells filled with 1:10 Mueller-Hinton Broth plus glyphosate at different concentrations served as negative controls. Positive controls were wells with L-asparagine and 1:10 Muller-Hinton Broth without glyphosate in which bacterial cells have grown as biofilm.


At the end of incubations, wells were aspirated and washed with 145 mM NaCl. Biofilms were fixed with 200 µL of 99% methanol (15 min). Methanol was removed and plates were air dried. Aliquots of 200 µL of 0.5% crystal violet (CV) were added to the wells. After 20 min, excess of CV was removed by washing with water. Finally, the impregnated CV was released by adding 33% acetic acid (150 µL) and the OD<sub>590nm</sub> was determined. Negative controls were wells with L-asparagine and 1:10 Muller-Hinton Broth plus glyphosate (at 0.845 ppm, 1.690 ppm, 8.45 ppm, 16.90 ppm, 84.50 ppm, 169 ppm, 845 ppm, and 1690 ppm) but without bacterial cells. Positive controls were wells washed with L-asparagine and filled with 1:10 Muller-Hinton Broth without glyphosate in which bacterial cells were challenged to form biofilms. Absorbance values of the blanks were subtracted from values obtained in treatments to eliminate spurious results due to background interference.

## Statistics

All tests described above were carried out in triplicate in at least three different situations. Numeric data were tabulated in MSEXcel® spreadsheets (Microsoft Co.). Obtained data were tested for normal distribution by the Kolmogorov-Smirnov test and for homogeneity of variance by Chi-square test. To determine the interference in biofilms formation, the statistical differences were assessed using Tukey HSD test. Comparisons in a same glyphosate concentration in different atmospheres were carried out by Tukey test. A threshold of 0.05 was used to determine differences. Statistical analyzes were conducted using the package BioEstat 5.0 (IDS Mamirauá).

## Results

The growth curves containing lag, log and early stationary phases were constructed with OD540nm values read at one-hour intervals, for a total period of twenty-four hours ([Figure 1](#)). Aerobic growth was superior to anaerobic one. In normoxia, lower glyphosate concentrations (0.845 ppm, 1.690 ppm, 8.45 ppm, 16.90 ppm) seemed to favor planktonic cell growth. The most interesting finding occurred in anoxic atmosphere, where positive-control cultures, although adapted to the lack of oxygen, did not show good growth and cultures in the presence of higher concentrations of glyphosate have markedly developed (cluster “c” in [Figure 1](#)).



[Figure 1](#)

Growth of *P. aeruginosa* ATCC®15442 in different concentrations of glyphosate and normoxic/anoxic conditions. Each point represents the arithmetical average of nine repetitions. Brackets arbitrarily group the final growths (*i.e.* after 24 h) which ...

Regarding the aerobic growth, from 14–16 h post-inoculation, there were detected two clusters. Cluster “a” comprises of glyphosate at concentrations below 16.9 ppm; on the other hand, cluster “b” comprises treatments with 84.5 ppm to 1690 ppm and the glyphosate-free control.

Biofilms of *P. aeruginosa* ATCC®15442 had more cells when were formed in absence of molecular oxygen ( $p < 0.05$ ); among the different treatments, the only exception occurred in 0.845 ppm glyphosate, where the means did not differ ( $p > 0.05$ ), although there seems to be a trend of increase in anoxia ([Figure 2](#)). Above 169 ppm, biofilms formed in anoxia retained more crystal violet, indicating higher cell content (Tukey HSD,  $p < 0.05$ ). The same occurred with biofilms formed in normoxia at concentrations above 845 ppm (Tukey HSD,  $p < 0.05$ ). A greater range of variation between biofilms formed in different atmospheres occurred at 1690 ppm (Student’s t test,  $p = 0.0003$ ).




Figure 2

Differential growth of 36 h-biofilms of *P. aeruginosa* ATCC®15442™ in different concentrations of glyphosate and in normoxic/anoxic conditions. Each bar represents the arithmetical average of nine repetitions with its respective 95% confidence ...

## Discussion

Glyphosate is probably the herbicide most discharged into the environment. Due to its extensive use in the protection of crops, it is inevitable that it will reach surface and deep waters ([Pournaras et al., 2007](#)), especially after rainfalls. The concentrations of glyphosate used herein are slightly higher than those found in surface waters after rainfalls ([Peruzzo et al., 2008](#)); however, as it is not possible to establish a consensus in terms of concentration, once it depends on variations in the amount applied, rainfall intensity, and degradation, we assumed that such dissonance should not be a major factor.

Aerobic planktonic growth was superior to anaerobic one. This points to the possibility of *P. aeruginosa*, although a facultative organism ([Davies et al., 1989](#); [Yoon et al., 2002](#)), has its growth significantly favored by the presence of molecular oxygen. Continuous bacterial exposure to low concentrations of glyphosate leads to increased rates of aerobic growth, which is somehow in agreement with previously published findings ([Fitzgibbon and Braymer, 1988](#)). By the contrary, in conditions of inaccessibility of molecular oxygen, the bacterium started to grow better in a concentration-dependent manner. It is possible that this phenomenon results from the use of the molecule as a source of phosphorus, as previously reported for the genus *Pseudomonas* ([Peñaloza-Vazquez et al., 1995](#); [Moore et al., 1983](#); [Talbot et al., 1984](#)). Glyphosate could also serve as a carbon source, which would be processed by both aerobic and anaerobic metabolisms ([Rueppel et al., 1977](#)), with increased rates in presence of oxygen. To support such theory, it has been found that different bacterial genera may promote catalysis of glyphosate using C-P lyases ([van Eerd et al., 2003](#)). Once broken up this connection, *Pseudomonas* spp. can produce glycine ([Kishore and Jacob, 1987](#)), which can also enhance growth.

To our knowledge, no other group has evaluated the effects that glyphosate can exert on sessile bacterial communities, so-called biofilms. Our results revealed that the xenobiotic tends to favor the formation of biofilms of *P. aeruginosa*, especially those anaerobic and that such increase seems to be concentration-dependent. Results of anaerobic planktonic and biofilm growth deserve greater reflection. In a hypothetical situation in which *P. aeruginosa* come to adhere to surfaces under low oxygen tensions and constant challenge of glyphosate, the growth can be incremented.

As Muller-Hinton broth, even ten-fold diluted, still is a culture medium with a complex composition and we may not ignore the possibility of some constituent had influenced the growth rates. However, this medium has been used in studies similar to this one, successfully ([Alandejani et al., 2009](#); [Pournaras et al., 2007](#)).

The results from this study point to the fact that the indiscriminate use of agricultural formulations containing glyphosate may result in an increase in growth rates of planktonic

and biofilm phenotypes of *P. aeruginosa* watercourses or reservoirs.

## Acknowledgments

The authors thank Professor Luiz Alberto Kozlowski (The Pontifical Catholic University of Paraná, Brazil), who kindly provided the glyphosate used in this study.

[Author information](#) ► [Article notes](#) ► [Copyright and License information](#) ►

## References

1. Alandejani T, Marsan J, Ferris W, Slinder R, Chan F. Effectiveness of honey on *Staphylococcus aureus* and *Pseudomonas aeruginosa* biofilms. *Otolaryngol Head Neck Surg.* 2009;141:114–118. [[PubMed](#)]
2. Araújo AS, Monteiro RT, Abarkeli RB. Effect of glyphosate on the microbial activity of two Brazilian soils. *Chemosphere.* 2003;52:799–804. [[PubMed](#)]
3. Dallegre E, Mantese FD, Oliveira RT, Andrade AJ, Dalsenter PR, Langeloh A. Pre and postnatal toxicity of the commercial glyphosate formulation in Wistar rats. *Arch Toxicol.* 2007;81:665–673. [[PubMed](#)]
4. Davies KJ, Lloyd D, Boddy L. The effect of oxygen on denitrification in *Paracoccus denitrificans* and *Pseudomonas aeruginosa*. *J Gen Microbiol.* 1989;135:2445–2451. [[PubMed](#)]
5. Fitzgibbon J, Braymer HD. Phosphate starvation induces uptake of glyphosate by *Pseudomonas* sp. strain PG2982. *Appl Environ Microbiol.* 1988;54:1886–1888. [[PMC free article](#)] [[PubMed](#)]
6. Govan JR, Deretic V. Microbial pathogenesis in cystic fibrosis: mucoid *Pseudomonas aeruginosa* and *Burkholderia cepacia*. *Microbiol Rev.* 1996;60:539–574. [[PMC free article](#)] [[PubMed](#)]
7. Kishore GM, Jacob GS. Degradation of glyphosate by *Pseudomonas* sp. PG2982 via a sarcosine intermediate. *J Biol Chem.* 1987;262:12164–12168. [[PubMed](#)]
8. Means NE, Kremer RJ, Ramsier C. Effects of glyphosate and foliar amendments on activity of microorganisms in the soybean rhizosphere. *J Environ Sci Health B.* 2007;42:125–132. [[PubMed](#)]
9. Moore JK, Braymer HD, Larson AD. Isolation of a *Pseudomonas* sp. which utilizes the phosphonate herbicide glyphosate. *Appl Environ Microbiol.* 1983;46:316–320. [[PMC free article](#)] [[PubMed](#)]
10. Peñaloza-Vazquez A, Mena GL, Herrera-Estrella L, Bailey AM. Cloning and sequencing of the genes involved in glyphosate utilization by *Pseudomonas pseudomallei*. *Appl Environ Microbiol.* 1995;61:538–543. [[PMC free article](#)] [[PubMed](#)]
11. Pérez GL, Torremorell A, Mugni H, Rodríguez P, Solange Vera M, do Nascimento M, Allende L, Bustingorry J, Escaray R, Ferraro M, Izaguirre I, Pizarro H, Bonetto C, Morris DP, Zagarese H. Effects of the herbicide Roundup on freshwater microbial communities: a mesocosm study. *Ecol Appl.* 2007;17:2310–2322. [[PubMed](#)]
12. Peruzzo PJ, Porta AA, Ronco AE. Levels of glyphosate in surface waters, sediments and soils

associated with direct sowing soybean cultivation in north pampasic region of Argentina. Environ Pollut. 2008;156:61–66. [[PubMed](#)]

13. Pournaras S, Ikonomidis A, Markogiannakis A, Spanakis N, Maniatis AN, Tsakris A. Characterization of clinical isolates of *Pseudomonas aeruginosa* heterogeneously resistant to carbapenems. J Med Microbiol. 2007;56:66–70. [[PubMed](#)]

14. Rueppel ML, Brightwell BB, Schaefer J, Marvel JT. Metabolism and degradation of glyphosphate in soil and water. J Agric Food Chem. 1977;25:517–528. [[PubMed](#)]

15. Stella J, Ryan M. Glyphosate herbicide formulation: a potentially lethal ingestion. Emerg Med Australas. 2004;16:235–239. [[PubMed](#)]

16. Talbot HW, Johnson LM, Munnecke DM. Glyphosate utilization by *Pseudomonas* sp. and *Alcaligenes* sp. isolated from environmental sources. Curr Microbiol. 1984;10:255–259.

17. Tsui MT, Chu LM. Comparative toxicity of glyphosate-based herbicides: aqueous and sediment porewater exposures. Arch Environ Contam Toxicol. 2004;46:316–323. [[PubMed](#)]

18. van Eerd LL, Hoagland RE, Zablotowicz RM, Hall JC. Pesticide metabolism in plants and microorganisms. Weed Sci. 2003;51:472–495.

19. Yoon SS, Hennigan RF, Hilliard GM, Ochsner UA, Parvatiyar K, Kamani MC, Allen HL, DeKievit TR, Gardner PR, Schwab U, Rowe JJ, Iglewski BH, McDermott TR, Mason RP, Wozniak DJ, Hancock RE, Parsek MR, Noah TL, Boucher RC, Hassett DJ. *Pseudomonas aeruginosa* anaerobic respiration in biofilms: relationships to cystic fibrosis pathogenesis. Dev Cell. 2002;3:593–603. [[PubMed](#)]

20. Wolfe RS. Techniques for cultivating methanogens. Methods Enzymol. 2011;494:1–22. [[PubMed](#)]

---

Articles from Brazilian Journal of Microbiology are provided here courtesy of Brazilian Society of Microbiology

The original source of this article is [Brazilian Journal of Microbiology](#)  
Copyright © [Global Research News](#), [Brazilian Journal of Microbiology](#), 2015

---

[Comment on Global Research Articles on our Facebook page](#)

[Become a Member of Global Research](#)

Articles by: [Global Research News](#)

**Disclaimer:** The contents of this article are of sole responsibility of the author(s). The Centre for Research on Globalization will not be responsible for any inaccurate or incorrect statement in this article. The Centre of Research on Globalization grants permission to cross-post Global Research articles on community internet sites as long the source and copyright are acknowledged together with a hyperlink to the original Global Research article. For publication of Global Research articles in print or other forms including commercial internet sites, contact: [publications@globalresearch.ca](mailto:publications@globalresearch.ca)  
[www.globalresearch.ca](http://www.globalresearch.ca) contains copyrighted material the use of which has not always been specifically authorized by the

copyright owner. We are making such material available to our readers under the provisions of "fair use" in an effort to advance a better understanding of political, economic and social issues. The material on this site is distributed without profit to those who have expressed a prior interest in receiving it for research and educational purposes. If you wish to use copyrighted material for purposes other than "fair use" you must request permission from the copyright owner.

For media inquiries: [publications@globalresearch.ca](mailto:publications@globalresearch.ca)