

Israel Reports Link Between Rare Cases of Heart Inflammation and COVID-19 Vaccination in Young Men

By [Gretchen Vogel](#) and [Jennifer Couzin-Frankel](#)

Global Research, June 06, 2021

[Science Magazine](#) 1 June 2021

Region: [Middle East & North Africa](#)

Theme: [Science and Medicine](#)

All Global Research articles can be read in 51 languages by activating the “Translate Website” drop down menu on the top banner of our home page (Desktop version).

Visit and follow us on Instagram at [@crg_globalresearch](#).

The COVID-19 vaccine made by Pfizer and BioNTech appears to put young men at elevated risk of developing a heart muscle inflammation called myocarditis, researchers in Israel say. In a report submitted today to the Israeli Ministry of Health, they conclude that between one in 3000 and one in 6000 men ages 16 to 24 who received the vaccine developed the rare condition. But most cases were mild and resolved within a few weeks, which is typical for myocarditis. “I can’t imagine it’s going to be anything that would cause medical people to say we shouldn’t vaccinate kids,” says Douglas Diekema, a pediatrician and bioethicist at Seattle Children’s Hospital.

Israeli health officials first flagged the issue in April, when they reported more than 60 cases, mostly in young men who had received their second dose of vaccine a few days earlier. Around the same time, the U.S. Department of Defense began [to track 14 such cases](#). In mid-May, the U.S. Centers for Disease Control and Prevention said [it, too, was reviewing myocarditis cases](#). Officials at the European Medicines Agency said on 28 May they had received 107 reports of myocarditis following the Pfizer-BioNTech vaccine, or about one in 175,000 doses administered. But relatively few people under age 30 have been vaccinated in Europe.

The Israeli panel’s findings come as Israel and many European countries are debating whether younger adolescents should be vaccinated against COVID-19. Israel has been vaccinating teenagers 16 and older since late January, and the Ministry of Health is scheduled to announce tomorrow whether vaccinations will be opened to children 12 and older. Other countries, including the United States and Canada, began vaccinating children 12 and older in mid-May.

“From a parent’s perspective, this really comes down to risk perception, assessment of the data,” says Diekema, who has studied risk-benefit trade-offs. Even if a link between myocarditis and the vaccine holds up, the condition is usually mild, requiring treatment only with anti-inflammatory drugs, whereas COVID-19 infection can also cause serious disease and long-term side effects, even in young people. As suspicion has percolated about a

possible connection, “I don’t know many physicians who are changing their minds about vaccinating their kids,” Diekema says.

In Israel, which relied almost exclusively on the Pfizer-BioNTech vaccine in its early and fast vaccination drive, the Ministry of Health in January assembled a panel led by Dror Mevorach, head of internal medicine at the Hadassah University Medical Center, to investigate the issue. Mevorach tells *Science* he and his colleagues identified 110 myocarditis cases among 5 million people in Israel who had received two doses of the Pfizer-BioNTech vaccine in the month before their diagnosis. That translates to about one in 50,000 vaccine recipients, a number that isn’t concerning given the background rate of myocarditis in the general population, where it is typically triggered by [viral or bacterial infections, including COVID-19](#).

But the rate of myocarditis following vaccination among young men was higher. Ninety percent of the cases picked up in Israel appeared in men, and although myocarditis is normally more common among young men, the rate among those vaccinated was somewhere between five and 25 times the background rate, the report says. (Two cases of fatal myocarditis have also been reported in Israel, but the panel says investigations of those deaths were inconclusive; one patient may have had a more generalized inflammatory syndrome, and the other diagnosis was “not verified,” the report says.)

The new analysis “is very suggestive of a causal nature,” between the vaccine and myocarditis, Mevorach says. “I am convinced there is a relationship.”

“It does suggest that this is, at least statistically, a real phenomenon,” says Peter Liu, a cardiologist and chief scientific officer of the University of Ottawa Heart Institute. Diekema says it’s important to investigate “even a hint of a signal,” but cautions that “while this report is suggestive ... it requires validation in other populations by other investigators before we can be certain the link exists.” Other factors may be in play, Diekema says. Now that children are back to socializing and playing sports, his hospital’s emergency room is “seeing more viral diseases than we’ve seen in a year,” and as a result, “I would expect to see a little bump in myocarditis versus a year ago.” Ideally, scientists should compare cohorts of vaccinated and unvaccinated youngsters at the same time, Diekema says, and he’s heartened that such studies are now gearing up.

Myocarditis cases following the Moderna vaccine, which isn’t in use in Israel, are also being investigated in the United States. It’s not clear why the two vaccines, which both rely on messenger RNA (mRNA), might heighten the risk. One possibility is that the very high antibody levels that both generate in young people may also, in rare cases, lead to a sort of immune overreaction that inflames the heart. “There’s no question these [vaccines] are extremely immune-generating,” Liu says. Mevorach says he suspects the mRNA itself might be playing a role. The innate immune system recognizes RNA as part of the body’s defense against microbes—including RNA viruses like SARS-CoV-2, he notes. “I think that actually the mRNA is a kind of natural adjuvant,” which ramps up the immune response, he says.

Diekema says the medical community is now on alert for youngsters with chest pain and other symptoms soon after vaccination—allowing them to be quickly identified, treated, and reported to health departments. Mevorach agrees that awareness among vaccinees, their parents, and their doctors is important for prompt and effective treatment. He says he and his colleagues treated about 40 cases. Only a few needed corticosteroids, he said, and most have recovered fully.

One important question is whether delaying the second vaccine dose might reduce any potential risk. There may be an opportunity to find out: Several countries have stretched the interval between the two doses from the 3 weeks tested and recommended by Pfizer to 12 or even 16 weeks, because they want to give as many people as possible at least one shot. A drop in myocarditis cases among those whose second dose was delayed might show up in data in the months ahead. Lowering the dose in young people may also be worth considering, Liu says. Pfizer's and Moderna's vaccines are now being tested at lower doses in children under 12, with results expected in the coming months.

Even if the link between the shots and myocarditis firms up, Liu says the vaccine's benefit—being well-protected from COVID-19—outweighs the risks, even for young people, who are generally at lower risk of severe disease. But Mevorach says the trade-offs may be different in Israel, given its extremely low numbers of SARS-CoV-2 infections—just 15 new cases were diagnosed yesterday. He hopes the Ministry of Health will leave the decision on whether to vaccinate younger teens to their parents and doctors. “At the moment, we no longer have an emergency,” he says.

Update, 1 June 2021, 4.55 PM: *Text has been added to this story about two reported fatal cases of myocarditis; the expert panel said investigations of those cases were inconclusive.*

*

Note to readers: Please click the share buttons above or below. Follow us on Instagram, @crg_globalresearch. Forward this article to your email lists. Crosspost on your blog site, internet forums. etc.

Gretchen Vogel is a contributing correspondent for *Science Magazine* based in Berlin, Germany.

[Jennifer Couzin-Frankel](#) is a Staff Writer.

The original source of this article is [Science Magazine](#)

Copyright © [Gretchen Vogel](#) and [Jennifer Couzin-Frankel](#), [Science Magazine](#), 2021

[Comment on Global Research Articles on our Facebook page](#)

[Become a Member of Global Research](#)

Articles by: **[Gretchen Vogel](#)**
and **[Jennifer Couzin-Frankel](#)**

Disclaimer: The contents of this article are of sole responsibility of the author(s). The Centre for Research on Globalization will not be responsible for any inaccurate or incorrect statement in this article. The Centre of Research on Globalization grants permission to cross-post Global Research articles on community internet sites as long the source and copyright are acknowledged together with a hyperlink to the original Global Research article. For publication of Global Research articles in print or other forms including commercial internet sites, contact: publications@globalresearch.ca

www.globalresearch.ca contains copyrighted material the use of which has not always been specifically authorized by the copyright owner. We are making such material available to our readers under the provisions of "fair use" in an effort to advance a better understanding of political, economic and social issues. The material on this site is distributed without profit to those who have expressed a prior interest in receiving it for research and educational purposes. If you wish to use copyrighted material for purposes other than "fair use" you must request permission from the copyright owner.

For media inquiries: publications@globalresearch.ca