

“It’s The Vaccines Stupid!”

Part I: Evidence Linking Autism Rise in Children to Vaccinations

By [F. William Engdahl](#)

Global Research, September 05, 2014

Global Research 5 September 2009

Theme: [Media Disinformation](#), [Science and Medicine](#)

In-depth Report: [THE H1N1 SWINE FLU PANDEMIC](#)

Following new revelations from U.S. Centers for Disease Control whistleblower William Thompson about mercury links to autism, we bring to our readers attention this important article first published by Global Research, 5 years ago today, September 5, 2009.

The WHO and US Government CDC are escalating a public psychological conditioning to create hysteria and panic among an uninformed public about an alleged “virus” H1N1 Influenza A, aka Swine Flu, whose alleged effects to date appear comparable with a common cold. Before people line up in the streets demanding their vaccinations for their children and themselves, it would be wise to remember, to paraphrase a 1992 campaign statement of Bill Clinton to George H.W. Bush: “It’s the vaccination, Stupid!”

By countless scientific accounts, far more dangerous to human health than any reported incidences of Swine Flu are the dangers of severe health issues including paralysis, brain damage and even death arising from what is added to vaccines by virtually every major vaccine maker. Almost without exception, all commercial vaccines today contain various substances known as adjuvants designed to make the vaccine “work.” These adjuvants are the source of horrendous and sometimes deadly damage.

It has been speculated for some time that there might be a link in the alarming rise in cases of autism among tiny infants and children and massive multiple vaccinations today given routinely to infants and children from the first hours of birth. There is clear and shocking evidence of the link between the two. If you do not have a strong constitution, you are advised not to read further.

A new study shows a direct link between standard childhood vaccination series, MMR, and autism-like symptoms in monkeys. The principal scientist involved in the study, Dr. Laura Hewitson of the University of Pittsburgh, presented the alarming conclusions as an abstract pending publication at the International Meeting for Autism Research. It has been presented at scientific conferences in both London and Seattle, USA.

The study compared vaccinated macaque monkeys with non-vaccinated macaques. No major flaws in the study have been revealed by any attending scientist. The vaccines included the popular MMR series. The study found a marked increase in “gastrointestinal tissue gene expression” and “inflammation issues” with those monkeys which received vaccinations. They are a common symptom of children with regressive autism.

The study also found marked behavior changes and development differences in those monkeys given the vaccines versus those who were not. “Compared with unexposed

animals, significant neuro-developmental deficits were evident for exposed animals in survival reflexes, tests of color discrimination and reversal, and learning sets,” the study’s authors reported. “Differences in behaviors were observed between exposed and unexposed animals and within the exposed group before and after MMR vaccination.”

US Government-mandated research approved by Congress was to begin this year, but the funds were rescinded in early January. Claiming “conflict of interest” because of ongoing court cases, the Centers for Disease Control and Prevention (CDC), a long-time supporter of infant vaccinations, withdrew the research plans.

The most shocking of all is the recent and now common medical practice, reinforced by an aggressive pharmaceutical industry, of giving multiple vaccines, often virtually within hours of birth, to infants despite the fact that no study including all of the vaccine series commonly given to children in the US and UK, about 30 in all, has been conducted until now. The practice of newborn multiple vaccinations has become widespread in Germany and other EU countries over the past decade. Significantly there have surfaced reports of dramatically increased instances of autism in newborn and infants in various German hospitals over the past decade, precisely the period multiple vaccinations of newborn and infants has become routine.

US Government coverup

Tragically, the US Government agency theoretically entrusted with guarding public health, the Food and Drug Administration (FDA), as with the case of health dangers of GMO foods, as well with the dramatic evidence of the link between autism and adjuvants used in typical vaccines, is accepting the argument of big and politically powerful Pharmaceutical companies.

The Food and Drug Administration considers vaccines safe but, just as with GMO, they have done no studies into the effects of multiple vaccinations as given in the common childhood series which started in the 1990s in the USA and spread to the UK and now across the EU.

According to Robert F. Kennedy, Jr., son of the late Attorney General and an attorney active in campaigning to expose mercury (Thimerosal) and other toxicity dangers in vaccines, recently stated, “as autism is a behavioral affliction rather than a precisely defined biological injury — epidemiological studies are critical to establishing its causation. But the greatest source of epidemiological data is the Vaccine Safety Datalink (VSD) — the government maintained medical records of hundreds of thousands of vaccinated children — which Health and Human Services Department has gone to great lengths to keep out of the hands of plaintiffs’ attorneys and independent scientists...The raw data collected in the VSD would undoubtedly provide the epidemiological evidence needed to understand the relationship between vaccines and autism. The absence of such studies makes it easy for judges to say to plaintiffs they have not met their burden of proving causation.”

Autism was virtually unknown in the United States until 1943 when it was diagnosed and identified eleven months after Thimerosal, a mercury-based vaccine “adjuvant” was first added to baby vaccines along with various aluminium compounds in the United States. Thimerosal is often used to stem fungi and bacterial growth in vaccines despite massive evidence of its severe effects as a potent neurotoxin. Following independent studies, Russia, Japan, Austria, Denmark, Sweden and Britain have banned Thimersol from childrens’

vaccines. Germany to date has no such ban. The toxin was developed in 1930 by Eli Lilly. Tragically in 1991, despite overwhelming evidence to the contrary the US Government's Center for Disease Control (CDC), the same agency fuelling the current hysteria over the non-proven H1N1 Swine Flu virus danger, recommended that infants be injected with a series of mercury-containing vaccines in some cases within 24 hours of birth for Hepatitis B and two months for diphtheria-tetanus-pertussis.

Before 1989 US pre-school children received eleven vaccinations—polio, diphtheria-tetanus-pertussis, measles-mumps-rubella (MMR). By 1999, because of the various CDC recommendations, the number of vaccinations was twenty two before first grade of school. Parallel with this explosive rise in vaccinations of the very young in the United States, according to Kennedy, the rate of autism among children. The state of Iowa reported a 700% increase in autism in children beginning in the 1990's and along with California has banned mercury in vaccines. Despite evidence, however the US FDA continues to allow drug makers to include Thimerosal in numerous over-the-counter non-prescription medications as well as steroids and injected collagen. The US Government ships vaccines preserved with Thimerosal to numerous developing countries as well, where some are reporting sudden explosion of autism rates as well. In China, where autism was unknown before introduction of Thimerosal by US drug makers in 1999, press reports indicate there are almost two million autistic children.

Instances of autism in the US exploded as some 40 million children were injected during the 1990's with Thimersol-based vaccines, giving them unprecedented accumulations of mercury poison. The level of ethylmercury in a vaccine routinely given then to children of two months age was 99 times greater than the US Government's daily limit for exposure. As with the current WHO pandemic declaration around H1N1 Swine Flu, the CDC Vaccine Advisory Committee is filled with scientists with close ties to the pharmaceutical industry. Dr. Sam Katz, chairman of the committee was a paid consultant to most companies producing the vaccines he "recommended."

The aluminium danger remains

While vaccines available in the US today exist with no Thimerosal (50% mercury), virtually all vaccines still contain aluminum, which has been linked to impaired neurological development in children. Aluminum has not replaced thimerosal as a vaccine preservative; it has always been used in vaccines.

In the recent past, most US children got exposed to both thimerosal and aluminum simultaneously with the hepatitis B, Hib, DTaP (diphtheria, tetanus and pertussis) and pneumococcal vaccines. Combining mercury with aluminum increases the likelihood that the mercury will damage human tissue.

According to a recent report by Michael Wagnitz, an American chemist, "Currently eight childhood vaccines that contain aluminum ranging from 125 to 850 micrograms (mcg). These vaccines are administered 17 times in the first 18 months of life, an almost six-fold increase compared to the vaccine schedule of the 1980s."

Wagnitz adds, "According to the American Society for Parenteral and Enteral Nutrition, based on IV feeding solutions, a child should not exceed a maximum daily dose of 5 mcg of aluminum per kilogram of weight per day. That means if a child weighs 11 pounds, the child should not exceed 25 mcg in a day. This level was determined to be the maximum safety

limit based on a study published in the New England Journal of Medicine titled “Aluminum Neurotoxicity in Preterm Infants Receiving Intravenous Feeding Solutions.”

The hepatitis B vaccine, administered at birth, contains 250 mcg.

In a 1996 policy statement, “Aluminum Toxicity in Infants and Children,” the American Academy of Pediatrics states, “Aluminum can cause neurological harm. People with kidney disease who build up bloodstream levels of aluminum greater than 100 mcg per liter are at risk of toxicity. The toxic threshold of aluminum in the bloodstream may be lower than 100 mcg per liter.” What level of aluminium toxicity is contained in vaccines routinely given German, French and other children in the EU is not known. It might be time for a public demand for such information to be disclosed, and before governments launch mass vaccination campaigns for untested vaccines against a non-proven H1N1 Swine Flu threat.

The original source of this article is Global Research
Copyright © [F. William Engdahl](#), Global Research, 2014

[Comment on Global Research Articles on our Facebook page](#)

[Become a Member of Global Research](#)

Articles by: [F. William Engdahl](#)

Disclaimer: The contents of this article are of sole responsibility of the author(s). The Centre for Research on Globalization will not be responsible for any inaccurate or incorrect statement in this article. The Centre of Research on Globalization grants permission to cross-post Global Research articles on community internet sites as long the source and copyright are acknowledged together with a hyperlink to the original Global Research article. For publication of Global Research articles in print or other forms including commercial internet sites, contact: publications@globalresearch.ca

www.globalresearch.ca contains copyrighted material the use of which has not always been specifically authorized by the copyright owner. We are making such material available to our readers under the provisions of "fair use" in an effort to advance a better understanding of political, economic and social issues. The material on this site is distributed without profit to those who have expressed a prior interest in receiving it for research and educational purposes. If you wish to use copyrighted material for purposes other than "fair use" you must request permission from the copyright owner.

For media inquiries: publications@globalresearch.ca