

Kids Who Received mRNA COVID Vaccines Had Altered Immune Systems a Year Later

By [Dr. Suzanne Burdick](#)

Theme: [Science and Medicine](#)

Global Research, August 20, 2024


[Children's Health Defense](#) 19 August 2024

Children ages 5-11 who received two doses of Pfizer's mRNA COVID-19 vaccine had heightened levels of a type of antibody suggestive of an altered immune system response one year after vaccination, a new [peer-reviewed study](#) revealed.

The team of German researchers, led by [Dr. Robin Kobbe](#) with the Institute for Infection Research and Vaccine Development at the University Medical Center Hamburg-Eppendorf in Germany, looked at blood samples of 14 healthy children the day the children received dose one of Pfizer's shot, one month afterward and one year after the children received dose two.


VACCINE REPORTS

Delayed Induction of Noninflammatory SARS-CoV-2 Spike-Specific IgG4 Antibodies Detected 1 Year After BNT162b2 Vaccination in Children

 Kobbe, Robin MD^{*,†,‡,§}; Rau, Cornelius MD, MPhil^{*,†}; Schulze-Sturm, Ulf MD[‡]; Stahl, Felix MD[§]; Fonseca-Brito, Luis[§]; Diemert, Anke MD^{¶,||}; Lütgehetmann, Marc MD^{**}; Addo, Marylyn M. MD^{*,†,‡,§}; Arck, Petra MD^{¶,||}; Weskamm, Leonie M. PhD^{*,†,‡,§}

[Author Information](#) 

The Pediatric Infectious Disease Journal ():10.1097/INF.0000000000004488, July 30, 2024. | DOI:

10.1097/INF.0000000000004488 

[OPEN](#) [SDC](#) [PAP](#)

 Metrics

Screenshot from [PIDJ](#)

A year after the second dose, they found increased levels of [IgG4](#) antibodies in the children's blood, suggesting that their immune system switched its type of immune system response.

IgG4 is one of the four [subclasses of immunoglobulin](#), or antibodies, produced by [plasma cells](#) in the blood.

While prior studies have found [elevated levels of IgG4 in adults](#) after repeated [mRNA](#)

[COVID-19 vaccination](#), Kobbe and his co-authors said their investigation is the first showing it happens in children, too.

The researchers wrote in their report published July 30 in [The Pediatric Infectious Disease Journal](#), “IgG4 responses should gain more attention in health and disease, especially in the context of mRNA vaccination.”

“Understanding the unusual mechanism triggering IgG4 production is crucial,” they added, “as more mRNA [vaccines](#) are currently under development and could hit the global market soon.”

Heightened IgG4 Indicative of IgG4-related Disease

[Brian Hooker, Ph.D.](#), chief scientific officer at [Children’s Health Defense](#) (CHD), told [The Defender](#) the study’s findings are very concerning because elevated IgG4 may be indicative of [IgG4-related disease](#) — a “multi-organ, fibro-inflammatory condition that usually involves the pancreas, kidneys or salivary glands but could involve any other organ.”

“Seventy to 80% of those with the disease have [elevated IgG4](#),” Hooker said. “Although IgG4-related disease is treatable, the underlying autoimmune conditions are often chronic and will require a lifetime of treatment.”

The disease could be autoimmune in origin due to molecular mimicry from the [COVID-19](#) vaccine, Hooker said. “It is also analogous to systemic sarcoidosis which is an inflammatory condition caused by an immune system exaggeration leading to granulomas.”

According to the Cleveland Clinic, [granulomas](#) are “clusters of white blood cells that ‘wall off’ bacteria, a foreign object or something else it thought was harmful from the rest of your body.” They most often form in the lungs, but can also form in the liver, kidney, skin or other areas of the body.

As [The Defender previously reported](#), the mRNA COVID-19 vaccine’s propensity to alter the immune system’s functioning in this way is something discussed in [Byram Bridle, Ph.D.](#), and [Dr. Harvey Risch](#)’s new book, “[Toxic Shot](#): Facing the Dangers of the COVID ‘Vaccines.’”

Bridle, a viral immunologist who wrote the book’s chapter on the vaccine’s “immunological harms,” did not respond to The Defender’s comment request on the German study. However, Risch — professor emeritus of epidemiology at the Yale School of Public Health — told The Defender in an earlier interview:

“After three to four doses of the vaccine, the antibody response of the immune system gets shifted from an IgG1 [[immunoglobulin type 1](#)] or 2 response, which are neutralization responses, to an [IgG4 response](#), which is a tolerance response.”

“[Tolerance](#)” describes how the immune system reduces its overreaction to certain pathogens, for example, those related to food or seasonal allergies. This dampening of immune system surveillance could potentially leave people more vulnerable to infections and other [health issues](#), including [cancer](#).

IgG4 Response May Reduce Body's Ability to Fight Cancer

When the immune system is dominated by IgG4 antibodies, the body may be less able to fight off cancer.

The authors of an April 24 review article published in [Nature Reviews Immunology](#) explained:

"IgG4 competes with other antibody (sub)classes for binding to tumour antigens and owing to its anti-inflammatory properties blocks the induction of antitumour immune responses ...

"In the absence of an immune response, tumour cells have increased ability to proliferate and metastasize, resulting in disease progression and decreased survival. Immune evasion through class-switching to IgG4 has been observed in patients with melanoma, cholangiocarcinoma, colon cancer, pancreatic cancer and glioblastoma."

A 2022 study found that individuals with IgG4-related disease appeared to have a [higher risk of cancer](#)— especially pancreatic cancer and lymphoma — compared with the general population.

NIAID-funded Researcher Acknowledges Increased IgG4 from mRNA COVID Shots

In a [commentary](#) published Feb. 7, 2023, in *Science Immunology*, [Shiv Pillai, M.D., Ph.D.](#), professor of Medicine and Health Sciences and Technology at Harvard Medical School, raised questions about how increased levels of IgG4 antibodies from mRNA COVID-19 vaccines may negatively impact the immune system.

Pillai is also a program director at an [Autoimmune Center of Excellence](#) at Massachusetts General Hospital, which is funded by the National Institutes of Allergy and Infectious Disease (NIAID). This year NIAID paid [more than \\$650,000](#) to fund a study he's leading on IgG4-related disease.

Pillai acknowledged that mRNA COVID-19 vaccines appear to produce increased IgG4 levels but said, "Accurately deciphering the negative consequences, if any, of increased IgG4 levels will be difficult."

Pillai emphasized that "innumerable large studies" have shown repeated mRNA COVID-19 vaccination has protected people from severe COVID-19 symptoms and hospitalization — although the citation number he listed for this statement failed to detail any studies or reviews.

The results of recent studies linking repeated mRNA vaccines with increased IgG4 levels "nonetheless" warrant doing clinical studies on the effectiveness of spreading out mRNA vaccine boosters to possibly once per year, he said.

Pillai added that another option would be to only use mRNA antigens in the first vaccine dose for its priming effect.

The Defender reached out to Kobbe for comment on the [study's](#) findings but did not receive a response by the deadline.

*

Click the share button below to email/forward this article to your friends and colleagues. Follow us on [Instagram](#) and [Twitter](#) and subscribe to our [Telegram Channel](#). Feel free to repost and share widely Global Research articles.

[One Month Before Global Research's Anniversary](#)

Suzanne Burdick, Ph.D., is a reporter and researcher for The Defender based in Fairfield, Iowa.

Featured image is from CHD

The original source of this article is [Children's Health Defense](#)
Copyright © [Dr. Suzanne Burdick](#), [Children's Health Defense](#), 2024

[Comment on Global Research Articles on our Facebook page](#)

[Become a Member of Global Research](#)

Articles by: [Dr. Suzanne Burdick](#)

Disclaimer: The contents of this article are of sole responsibility of the author(s). The Centre for Research on Globalization will not be responsible for any inaccurate or incorrect statement in this article. The Centre of Research on Globalization grants permission to cross-post Global Research articles on community internet sites as long the source and copyright are acknowledged together with a hyperlink to the original Global Research article. For publication of Global Research articles in print or other forms including commercial internet sites, contact: publications@globalresearch.ca
www.globalresearch.ca contains copyrighted material the use of which has not always been specifically authorized by the copyright owner. We are making such material available to our readers under the provisions of "fair use" in an effort to advance a better understanding of political, economic and social issues. The material on this site is distributed without profit to those who have expressed a prior interest in receiving it for research and educational purposes. If you wish to use copyrighted material for purposes other than "fair use" you must request permission from the copyright owner.

For media inquiries: publications@globalresearch.ca