

Lab Rat Offspring Got Rib Malformations After COVID Vaccination: Moderna Trial Documents

By [Enrico Trigoso](#)

Global Research, August 18, 2022

[The Epoch Times](#) 16 August 2022

Region: [USA](#)

Theme: [Science and Medicine](#)

All Global Research articles can be read in 51 languages by activating the “Translate Website” drop down menu on the top banner of our home page (Desktop version).

To receive Global Research’s Daily Newsletter (selected articles), [click here](#).

Follow us on [Instagram](#) and [Twitter](#) and subscribe to our [Telegram Channel](#). Feel free to repost and share widely Global Research articles.

[Moderna](#) documents regarding their COVID vaccine trial on animals, obtained via a Freedom of Information Act request by Judicial Watch, showed that some of the offspring of [rats](#) that were injected with Moderna’s mRNA shot developed rib malformations.

The 700 pages contain a portion of the formal Biologics Licensing Application (BLA) package that a manufacturer is required to submit to the FDA for approval.

The documents have not yet been made public, but were analyzed by former pharma executive Alexandra Latypova and reviewed by The Epoch Times.

Included in the documents are test results that show that Moderna mRNA shots caused statistically significant skeletal malformations in the offspring of the rats that took the mRNA-1273 (Spikevax mRNA) doses.

“mRNA-1273-related variations in skeletal examination included statistically significant increases in the number of F1 rats with 1 or more wavy ribs and 1 or more rib nodules. Wavy ribs appeared in 6 fetuses and 4 litters with a fetal prevalence of 4.03% and a litter prevalence of 18.2%. Rib nodules appeared in 5 of those 6 fetuses,” according to Moderna’s internal documents.

F1 refers to the rat offspring and litter indicates a group-birth of rats.

“Maternal toxicity in the form of clinical observations was observed for 5 days following the last dose (Gestation Day 13), correlating with the most sensitive period for rib development in rats (Gestation Days 14 to 17)” the documents state.

In the F₁ generation, there were no mRNA-1273-related effects or changes in the following parameters: mortality, body weight, clinical observations, macroscopic observations, gross pathology, external or visceral malformations or variations, skeletal malformations, and mean number of ossification sites per fetus per litter. mRNA-1273-related variations in skeletal examination included statistically significant increases in the number of F₁ rats with 1 or more wavy ribs and 1 or more rib nodules. Wavy ribs appeared in 6 fetuses and 4 litters with a fetal prevalence of 4.03% and a litter prevalence of 18.2%. Rib nodules appeared in 5 of those 6 fetuses. Skeletal variations are structural changes that do not impact development or function of a developing embryo, are considered reversible, and often correlate with maternal toxicity and/or lack of other indicators of developmental toxicity (Carney and Kimmel 2007). Maternal toxicity in the form of clinical observations was observed for 5 days following the last dose (GD 13), correlating with the most sensitive period for rib development in rats (GDs 14 to 17). Furthermore, there were no other indicators of mRNA-1273-related developmental toxicity observed, including delayed ossification; therefore, these common skeletal variations were not considered adverse.

Overall, maternal administration of mRNA-1273 on Study Day 1 (28 days prior to mating), Study Day 15 (14 days prior to mating) and GD 1, and GD 13 did not have any adverse effects on the F₀ or F₁ generations. The mRNA-1273-related, non-adverse effects were limited to an increase in the number of fetuses with common skeletal variations of 1 or more rib nodules and 1

Confidential

Page 22
FDA-CBER-2021-4379-0001151

A document package on biodistribution studies obtained by Judicial Watch. (Courtesy of Alexandra Latypova)

“Wavy ribs” refers to ribs not properly shaped.

In other words, 6 out of about 149 baby rats had wavy ribs and 5 of those also had rib nodules.

According to Latypova’s analysis, only female rats were studied (male rats were not treated with the Moderna vaccine).

The females got a human dose of 100mcg Spikevax mRNA, 28 and 14 days prior to mating and gestation days 1 and 13.

“1/2 rats euthanized before delivery to examine fetuses, the rest followed to 21 days after delivery,” Latypova stated, “No numbers are provided for the study size.”

In addition, there is no study report, but only Moderna’s own interpretation of the outcomes.

The results were part of reproductive toxicology tests done by Moderna, which is the only reproductive toxicology test for the product, according to Latypova.

It is not known how the dose translates from humans to rats.

“Neither Moderna nor Pfizer provided any dose calculations or justification information for

dose selection in animal studies,” Latypova told The Epoch Times.

“Doses of drugs, or especially biologics do not necessarily have linear relationships with toxicity or efficacy. It is likely a much more complex relationship and unfortunately not known at all.”

“Reproductive toxicology is the study of adverse effects of medicinal products on reproduction. The FDA requires reproductive toxicity testing for any NME to be used in women of childbearing potential,” added Latypova, who has worked in more than 60 pharmaceutical companies, mainly focusing on creating and reviewing clinical trials, many of which were submitted to the FDA.

Despite the abovementioned lab results, the [FDA issued a statement](#) on Jan. 30 saying that there were no adverse effects on postnatal developments.

“No vaccine-related fetal malformations or variations and no adverse effect on postnatal development were observed in the study,” the FDA stated on the label for Moderna’s Spikevax vaccine.

“In a developmental toxicity study, 0.2 mL of a vaccine formulation containing nucleoside-modified mRNA (100 mcg) and other ingredients that are included in a 0.5-mL single human dose of SPIKEVAX was administered IM to female rats on four occasions: 28 and 14 days prior to mating, and on gestation days 1 and 13,” reads the FDA publication.

Pfizer Vaccine Also Caused Abnormal Ribs in Rats

In August of 2021, Elsevier published a peer-reviewed study ([pdf](#)) titled “Lack of effects on female fertility and prenatal and postnatal offspring development in rats with BNT162b2, a mRNA-based COVID-19 vaccine.”

BNT162b2 is the Pfizer jab.

All the authors of the publication were employed by Pfizer, BioNTech, or Charles River, a Pfizer contractor.

This publication suggests that there was a “lack of effects” in postnatal offspring development, but the study shows that there was a 295 percent increase (8.3 percent compared to 2.1 percent in the control group) in abnormal ribs in vaccinated rat offspring. A huge increase in what is described as the “supernumerary lumbar.”

‘This Was an Extremely Dangerous Warning’

“Skeletal abnormalities in the bony rib cages are absolutely important and were statistically increased in the rat offspring of the experimental group compared with the placebo group,” James Thorp, an MD board-certified in obstetrics and gynecology, as well as maternal-fetal medicine, told The Epoch Times after reviewing Latypova’s analysis of Moderna’s BLA package.

“In clinical obstetrics and maternal-fetal medicine we see similar findings in skeletal abnormalities prior to birth that are extremely serious and often lethal. This was an

extremely dangerous warning signal in reproductive toxicology studies and was never brought to the light of day to protect our global citizens. The CDC, Pfizer, Moderna, and the flagship medical journals of the medical industrial complex lied to the American public and should be held accountable,” Thorp said.

Thorp [recently analyzed](#) and verified the most recent Vaccine Adverse Event Reporting System (VAERS) data related to COVID-19 vaccines and compared them to the influenza vaccines, finding numerous abnormalities.

The [CDC website](#) recommends the COVID vaccines during pregnancy in order to “prevent severe illness and death in pregnant women.”

The [American College of Obstetricians and Gynecologists](#) also “strongly recommends that pregnant individuals be vaccinated against COVID-19,” adding that pregnant women’s complete vaccination should be a “priority.”

Moderna, Pfizer, the CDC, and the FDA did not respond to requests for comment.

*

Note to readers: Please click the share buttons above or below. Follow us on Instagram and Twitter and subscribe to our Telegram Channel. Feel free to repost and share widely Global Research articles.

Enrico Trigoso is an Epoch Times reporter focusing on the NYC area.

Featured image by Janet Stephens, licensed under the [Public Domain](#)

The original source of this article is [The Epoch Times](#)
Copyright © [Enrico Trigoso](#), [The Epoch Times](#), 2022

[Comment on Global Research Articles on our Facebook page](#)

[Become a Member of Global Research](#)

Articles by: [Enrico Trigoso](#)

Disclaimer: The contents of this article are of sole responsibility of the author(s). The Centre for Research on Globalization will not be responsible for any inaccurate or incorrect statement in this article. The Centre of Research on Globalization grants permission to cross-post Global Research articles on community internet sites as long the source and copyright are acknowledged together with a hyperlink to the original Global Research article. For publication of Global Research articles in print or other forms including commercial internet sites, contact: publications@globalresearch.ca

www.globalresearch.ca contains copyrighted material the use of which has not always been specifically authorized by the copyright owner. We are making such material available to our readers under the provisions of "fair use" in an effort to advance a better understanding of political, economic and social issues. The material on this site is distributed without profit to those who have expressed a prior interest in receiving it for research and educational purposes. If you wish to use copyrighted material for purposes other than "fair use" you must request permission from the copyright owner.

For media inquiries: publications@globalresearch.ca