

# Medical Journal Article: 14,000 U.S. Deaths Tied to Fukushima Reactor Disaster Fallout

By [Global Research](#)

Global Research, December 20, 2011

[PRNewswire-USNewswire](#) 20 December 2011

Region: [USA](#)

Theme: [Environment](#), [Science and Medicine](#)

## ***Impact Seen As Roughly Comparable to Radiation-Related Deaths After Chernobyl; Infants Are Hardest Hit, With Continuing Research Showing Even Higher Possible Death Count.***

WASHINGTON, Dec. 19, 2011 /PRNewswire-USNewswire/ — An estimated 14,000 excess deaths in the United States are linked to the radioactive fallout from the disaster at the Fukushima nuclear reactors in Japan, according to a major new article in the December 2011 edition of the International Journal of Health Services. This is the first peer-reviewed study published in a medical journal documenting the health hazards of Fukushima.

### [TO READ THE COMPLETE ARTICLE,](#)

*International Journal of Health Services*, Volume 42, Number 1, Pages 47-64, 2012 (based at the Johns Hopkins University, School of Hygiene and Public Health) [Global Research Editor's Note: IJHS is a prestigious peer reviewed journal.]

Authors Joseph Mangano and Janette Sherman note that their estimate of 14,000 excess U.S. deaths in the 14 weeks after the Fukushima meltdowns is comparable to the 16,500 excess deaths in the 17 weeks after the Chernobyl meltdown in 1986. The rise in reported deaths after Fukushima was largest among U.S. infants under age one. The 2010-2011 increase for infant deaths in the spring was 1.8 percent, compared to a decrease of 8.37 percent in the preceding 14 weeks.

The IJHS article will be published Tuesday and will be available online as of 11 a.m. EST at <http://www.radiation.org>.

Just six days after the disastrous meltdowns struck four reactors at Fukushima on March 11, scientists detected the plume of toxic fallout had arrived over American shores. Subsequent measurements by the U.S. Environmental Protection Agency (EPA) found levels of radiation in air, water, and milk hundreds of times above normal across the U.S. The highest detected levels of Iodine-131 in precipitation in the U.S. were as follows (normal is about 2 picocuries I-131 per liter of water): Boise, ID (390); Kansas City (200); Salt Lake City (190); Jacksonville, FL (150); Olympia, WA (125); and Boston, MA (92).

Epidemiologist Joseph Mangano, MPH MBA, said: **“This study of Fukushima health hazards is the first to be published in a scientific journal. It raises concerns, and strongly suggests that health studies continue, to understand the true impact of**

**Fukushima in Japan and around the world. Findings are important to the current debate of whether to build new reactors, and how long to keep aging ones in operation.”**

Mangano is executive director, Radiation and Public Health Project, and the author of 27 peer-reviewed medical journal articles and letters.

Internist and toxicologist Janette Sherman, MD, said: **“Based on our continuing research, the actual death count here may be as high as 18,000, with influenza and pneumonia, which were up five-fold in the period in question as a cause of death. Deaths are seen across all ages, but we continue to find that infants are hardest hit because their tissues are rapidly multiplying, they have undeveloped immune systems, and the doses of radioisotopes are proportionally greater than for adults.”**

Dr. Sherman is an adjunct professor, Western Michigan University, and contributing editor of *“Chernobyl – Consequences of the Catastrophe for People and the Environment”* published by the NY Academy of Sciences in 2009, and author of *“Chemical Exposure and Disease and Life’s Delicate Balance – Causes and Prevention of Breast Cancer.”*

The Centers for Disease Control and Prevention (CDC) issues weekly reports on numbers of deaths for 122 U.S. cities with a population over 100,000, or about 25-30 percent of the U.S. In the 14 weeks after Fukushima fallout arrived in the U.S. (March 20 to June 25), deaths reported to the CDC rose 4.46 percent from the same period in 2010, compared to just 2.34 percent in the 14 weeks prior. Estimated excess deaths during this period for the entire U.S. are about 14,000.

The original source of this article is [PRNewswire-USNewswire](#)  
Copyright © [Global Research](#), [PRNewswire-USNewswire](#), 2011

---

[Comment on Global Research Articles on our Facebook page](#)

[Become a Member of Global Research](#)

Articles by: [Global Research](#)

**Disclaimer:** The contents of this article are of sole responsibility of the author(s). The Centre for Research on Globalization will not be responsible for any inaccurate or incorrect statement in this article. The Centre of Research on Globalization grants permission to cross-post Global Research articles on community internet sites as long the source and copyright are acknowledged together with a hyperlink to the original Global Research article. For publication of Global Research articles in print or other forms including commercial internet sites, contact: [publications@globalresearch.ca](mailto:publications@globalresearch.ca)

[www.globalresearch.ca](http://www.globalresearch.ca) contains copyrighted material the use of which has not always been specifically authorized by the copyright owner. We are making such material available to our readers under the provisions of "fair use" in an effort to advance a better understanding of political, economic and social issues. The material on this site is distributed without profit to those who have expressed a prior interest in receiving it for research and educational purposes. If you wish to use copyrighted material for purposes other than "fair use" you must request permission from the copyright owner.

For media inquiries: [publications@globalresearch.ca](mailto:publications@globalresearch.ca)