

## 'Scale of this Failure Has No Precedent': Scientists Say Hot Ocean 'Blob' Killed One Million Seabirds

The lead author called the mass die-off "a red-flag warning about the tremendous impact sustained ocean warming can have on the marine ecosystem."

By [Jessica Corbett](#)

Theme: [Environment](#)

Global Research, January 20, 2020

[Common Dreams](#) 16 January 2020

*On the heels of [new research](#) showing that the world's oceans are rapidly warming, scientists [revealed](#) Wednesday that a huge patch of hot water in the northeast Pacific Ocean dubbed "the blob" was to blame for killing about one million seabirds.*

The peer-reviewed study, published in the journal *PLOS ONE*, was conducted by a team of researchers at federal and state agencies, conservation groups, and universities. They tied the mass die-off to "the blob," a marine heatwave that began forming in 2013 and grew more intense in 2015 because of the weather phenomenon known as El Niño.

"About 62,000 dead or dying common murres (*Uria aalge*), the tropically dominant fish-eating seabird of the North Pacific, washed ashore between summer 2015 and spring 2016 on beaches from California to Alaska," the study says. "Most birds were severely emaciated and, so far, no evidence for anything other than starvation was found to explain this mass mortality. Three-quarters of murres were found in the Gulf of Alaska and the remainder along the West Coast."

Given that previous studies have shown "that only a fraction of birds that die at sea typically wash ashore," the researchers put the death toll closer to a million.

"The magnitude and scale of this failure has no precedent," lead author John Piatt, a research biologist at the U.S. Geological Survey's Alaska Science Center and an affiliate professor at the University of Washington, [said](#) in a statement. "It was astonishing and alarming, and a red-flag warning about the tremendous impact sustained ocean warming can have on the marine ecosystem."

A New Study About the Death of 1 Million Seabirds Should Scare the Crap Out of You [#ClimateReality](#) [#ClimateAction](#)<https://t.co/7A17ZEGV1x>  
[pic.twitter.com/OJ01VIGlxP](https://pic.twitter.com/OJ01VIGlxP)

— CenterForTheBlueEconomy\_MIIS (@CBE\_MIIS) [January 16, 2020](#)

Piatt and study co-author and University of Washington professor Julia Parrish explained that

the team believes the blob—which spanned hundreds of miles—limited food supply in the region, leading the birds to starve.

“Think of it as a run on the grocery stores at the same time that the delivery trucks to the stores stopped coming so often,” Parrish said. “We believe that the smoking gun for common murre—beyond the marine heatwave itself—was an ecosystem squeeze: fewer forage fish and smaller prey in general, at the same time that competition from big fish predators like walleye, pollock, and Pacific cod greatly increased.”

Piatt added that

“food demands of large commercial groundfish like cod, pollock, halibut, and hake were predicted to increase dramatically with the level of warming observed with the blob, and since they eat many of the same prey as murre, this competition likely compounded the food supply problem for murre, leading to mass mortality events from starvation.”

According to *CNN*, which [reported](#) on the study Thursday:

The blob devastated the murre’s population. With insufficient food, breeding colonies across the entire region had reproductive difficulties for years afterward, the study said. Not only did the population decline dramatically, but the murre couldn’t replenish those numbers.

During the 2015 breeding season, three colonies didn’t produce a single chick. That number went up to 12 colonies in the 2016 season—and in reality it could be even higher, since researchers only monitor a quarter of all colonies.

Thomas Frölicher, a climate scientist at the University of Bern in Switzerland who was not involved in the new study, [discussed](#) the blob’s connection to the human-caused planetary emergency with *InsideClimate News*.

“It was the biggest marine heatwave so far on record,” said Frölicher, who noted that such events have doubled in frequency over the past few decades. “Usually, we are used to heatwaves over land. They are much smaller in size, and they do not last as long. In the ocean, this heatwave lasted two or three years.”

Frölicher warned that “if we follow a high-greenhouse-gas-emissions scenario, these heatwaves will become [50 times more frequent](#) than preindustrial times” by 2100. He said that even if the international community achieves a low-emissions scenario in line with the Paris climate agreement, marine heatwaves would still be 20 times more frequent.

“What that means is that in some regions, they will become permanent heatwaves,” he added. “This gives us some insight into the future.”

A severe marine heatwave in the Pacific Ocean killed almost a million seabirds. Our survival depends on drawing down greenhouse gas emissions.  
<https://t.co/F9r9xBtlyr>

The study—which its authors expect to inform research on other mortality events related to marine heatwaves—was published just weeks after University of Washington scientists found what [some have called](#) “the blob 2.0” forming in the Pacific. That discovery came as “quite a surprise” to those researchers.

University climatologist Nick Bond [told](#) local media that “the original blob was so unusual, and stood above the usually kind of variations in the climate and ocean temperatures, that we thought ‘wow, this is going to be something we won’t see for quite a while.’”

*From Common Dreams: Our work is licensed under a Creative Commons Attribution-Share Alike 3.0 License. Feel free to republish and share widely.*

\*

Note to readers: please click the share buttons above or below. Forward this article to your email lists. Crosspost on your blog site, internet forums. etc.

*Featured image: Dead common murres were found on the beach in Cochrane Bay, Prince William Sound on Jan 10, 2016. These birds were part of the large die-off of common murres across the Gulf of Alaska in 2015-2016. (Photo: Sarah Schoen/USGS Alaska Science Center)*

The original source of this article is [Common Dreams](#)  
Copyright © [Jessica Corbett](#), [Common Dreams](#), 2020

---

[Comment on Global Research Articles on our Facebook page](#)

[Become a Member of Global Research](#)

Articles by: [Jessica Corbett](#)

**Disclaimer:** The contents of this article are of sole responsibility of the author(s). The Centre for Research on Globalization will not be responsible for any inaccurate or incorrect statement in this article. The Centre of Research on Globalization grants permission to cross-post Global Research articles on community internet sites as long the source and copyright are acknowledged together with a hyperlink to the original Global Research article. For publication of Global Research articles in print or other forms including commercial internet sites, contact: [publications@globalresearch.ca](mailto:publications@globalresearch.ca)  
[www.globalresearch.ca](http://www.globalresearch.ca) contains copyrighted material the use of which has not always been specifically authorized by the copyright owner. We are making such material available to our readers under the provisions of "fair use" in an effort to advance a better understanding of political, economic and social issues. The material on this site is distributed without profit to those who have expressed a prior interest in receiving it for research and educational purposes. If you wish to use copyrighted material for purposes other than "fair use" you must request permission from the copyright owner.

For media inquiries: [publications@globalresearch.ca](mailto:publications@globalresearch.ca)